

## PLACENTAL BLOOD AS A SOURCE OF HEMATOPOIETIC STEM CELLS FOR TRANSPLANTATION INTO UNRELATED RECIPIENTS

JOANNE KURTZBERG, M.D., MARY LAUGHLIN, M.D., MICHAEL L. GRAHAM, M.D., CLAY SMITH, M.D.,  
JANICE F. OLSON, M.D., EDWARD C. HALPERIN, M.D., GILBERT CIOCCI, P.N.P.,  
CARMELITA CARRIER, PH.D., CLADD E. STEVENS, M.D., AND PABLO RUBINSTEIN, M.D.

### ABSTRACT

**Background** Transplantation of bone marrow from unrelated donors is limited by a lack of HLA-matched donors and the risk of graft-versus-host disease (GVHD). Placental blood from sibling donors can reconstitute hematopoiesis. We report preliminary results of transplantation using partially HLA-mismatched placental blood from unrelated donors.

**Methods** Twenty-five consecutive patients, primarily children, with a variety of malignant and nonmalignant conditions received placental blood from unrelated donors and were evaluated for hematologic and immunologic reconstitution and GVHD. HLA matching was performed before transplantation by serologic typing for class I HLA antigens and low-resolution molecular typing for class II HLA alleles. In donor-recipient pairs who differed by no more than one HLA antigen or allele, high-resolution class II HLA typing was done retrospectively. For donor-recipient pairs who were mismatched for two HLA antigens or alleles, high-resolution typing was used prospectively to select the best match for HLA-DRB1.

**Results** Twenty-four of the 25 donor-recipient pairs were discordant for one to three HLA antigens. In 23 of the 25 transplant recipients, the infused hematopoietic stem cells engrafted. Acute grade III GVHD occurred in 2 of the 21 patients who could be evaluated, and 2 patients had chronic GVHD. In vitro proliferative responses of T cells and B cells to plant mitogens were detected 60 days after transplantation. With a median follow-up of 12½ months and a minimal follow-up of 100 days, the overall 100-day survival rate among these patients was 64 percent, and the overall event-free survival was 48 percent.

**Conclusions** HLA-mismatched placental blood from unrelated donors is an alternative source of stem cells for hematopoietic reconstitution in children. (N Engl J Med 1996;335:157-66.)

©1996, Massachusetts Medical Society.

**A**LLOGENEIC bone marrow transplantation can cure some hematologic cancers, bone marrow failure syndromes, immunodeficiency disorders, and inborn errors of metabolism,<sup>1-5</sup> but its success depends on the prompt identification of a suitable donor and the avoidance of severe graft-versus-host disease (GVHD).<sup>6-10</sup> Transplantation of hematopoietic stem cells from placental or cord blood can overcome these problems.

Over the past seven years, placental blood from a

sibling has been used as a source of hematopoietic stem cells in more than 100 allogeneic transplantations.<sup>11-14</sup> In 44 placental-blood transplantations involving sibling donors that were reported to the International Cord Blood Transplant Registry, there was successful hematopoietic reconstitution and a lower incidence of GVHD than expected with bone marrow grafts.<sup>11,12</sup> Delays in myeloid engraftment were noted, but the probability of event-free survival was 72 percent after a median follow-up of 1.6 years.

In 1992 the Placental Blood Program was established at the New York Blood Center to explore the feasibility of using banked placental blood from unrelated donors for the transplantation of hematopoietic stem cells.<sup>15</sup> In this report we describe the preliminary results of 25 consecutive transplantations of placental blood from unrelated donors that were performed at a single center with units obtained through the Placental Blood Program. Nearly all of the patients in this study were children (age range, 0.8 to 23.5 years).

### METHODS

#### Eligibility and Study Objectives

This phase I study of the transplantation of placental blood for the treatment of malignant and nonmalignant conditions was approved by the institutional review board of Duke University Medical Center. Patients were eligible for enrollment if there was neither an HLA-identical related donor nor a related donor with two HLA mismatches available and if an HLA-matched, unrelated bone marrow donor could not be identified within six months. Informed consent and a sample of autologous backup bone marrow were also required.

#### Donor Selection

Beginning in September 1993, formal searches were conducted by the Placental Blood Program on behalf of 210 patients referred to the Duke University Medical Center. Initial matching criteria for an HLA-matched unit required the placental blood to share at least five HLA antigens with the potential recipient. These antigens were identified by serologic typing (HLA class I) and low-resolution DNA typing (HLA class II). Such matched units were found for 92 of the 210 patients (44 percent). Of these 92 units, 6 matched all six of the recipients' class I and class

From the Pediatric Bone Marrow Transplant Program at Duke University Medical Center, Durham, N.C. (J.K., M.L., M.L.G., C.S., J.F.O., E.C.H., G.C.), and the Placental Blood Program at the New York Blood Center, New York (C.C., C.E.S., P.R.). Address reprint requests to Dr. Kurtzberg at Box 3350, Duke University Medical Center, Durham, NC 27710.

II major histocompatibility complex (MHC) antigens. Donor units that matched with four of the prospective recipients' six class I and II HLA antigens were sought for 58 of the remaining patients and identified for 52.

When several units were available for a patient, we selected the one that best matched the patient's HLA haplotypes and contained an optimal number of nucleated cells. In choosing among units with one HLA incompatibility, mismatches at class II MHC loci were considered as important as class I mismatches; when possible, mismatches within the same cross-reacting group of HLA antigens were preferred to major incompatibilities. When the donor unit and the potential recipient differed by two HLA antigens, more than one unit was usually available. In these instances, the results of high-resolution typing of HLA-DRB1 were used to find the unit with the closest match. In all other cases, high-resolution typing was analyzed retrospectively. After identification of a suitable placental blood unit, confirmatory HLA typing of the patient and the prospective unit was performed. A test sample from the unit that was transplanted into the one patient with Lesch-Nyhan disease was assayed for hypoxanthine phosphoribosyltransferase activity by Dr. William Nyhan. None of the transplanted placental-blood units were depleted of red cells, reduced in volume, or depleted of T cells before cryopreservation. All transplanted units were negative for human immunodeficiency virus (HIV), hepatitis A, B, and C, and human T-cell lymphotropic

virus type I (HTLV-I) by standard blood-bank screening tests. None of the units, nor the mothers of the donors, were positive for IgM antibody to cytomegalovirus (CMV).

**Transplantation Procedure**

Cryopreserved units of placental blood were transported to the study center in a shipping container cooled by liquid nitrogen in vapor phase and stored in liquid nitrogen. For the first three transplantations, the unit of blood was thawed at the bedside and infused intravenously over a period of 10 to 15 minutes. For subsequent transplantations, the unit was thawed in the laboratory and washed with 10 percent dextran 40 (Baxter, Glendale, Calif.) and 5 percent human albumin before infusion.<sup>16</sup> A nucleated-cell count, ABO and Rh typing, a test of cell viability, bacterial and fungal cultures, an assay for hematopoietic progenitor cells, and a CD34+ cell count were performed on a sample from each thawed unit at the time of infusion (Table 1).

**Preparative Regimens and GVHD**

Table 2 lists the preparative regimens and details of prophylaxis against GVHD. Patients two years of age or older who had leukemia received hyperfractionated total-body irradiation (150 cGy twice a day for nine doses), melphalan, and antithymocyte globu-

**TABLE 1. CHARACTERISTICS OF PLACENTAL-BLOOD GRAFTS.**

PATIENT No.	ABO/RH TYPE	VOLUME COLLECTED	VOLUME INFUSED	No. OF	No. OF	No. OF COLONY-	No. OF
				NUCLEATED CELLS CRYOPRESERVED*	NUCLEATED CELLS INFUSED	FORMING UNITS INFUSED	CD34+ CELLS INFUSED
	patient/donor	ml		×10 <sup>-7</sup> /kg		×10 <sup>-4</sup> /kg	×10 <sup>-6</sup> /kg
1	O+/O+	98	148	4.6	2.7	2.16	3.65
2	O+/O-	48	123	4.8	3.7	6.29	ND†
3	A-/A+	44	108	0.7	0.7	0.63	ND†
4	A+/O+	88	155	6.0	6.8	22.40	1.36
5	A-/O+	48	152	2.8	2.1	4.41	2.10
6	A+/O+	91	90	5.4	4.8	20.00	1.35
7	B+/O+	63	125	1.6	1.3	2.34	0.70
8	A+/A-	53	72	2.7	2.1	3.60	0.44
9	A+/O+	43	51	2.1	2.4	3.80	0.63
10	A+/A+	65	111	2.0	1.5	2.70	0.38
11	A+/A+	59	93	5.5	2.0	4.20	0.17
12	B+/A+	51	75	2.3	1.6	1.30	0.05
13	O+/B+	128	129	6.3	3.7	3.80	2.90
14	A+/O+	68	70	9.0	5.4	3.20	8.93
15	A-/B+	63	81	2.6	3.0	9.90	1.43
16	B+/O+	104	122	4.2	3.8	0.23	0.72
17	A+/AB-	52	55	4.1	2.4	3.10	1.92
18	A+/A+	214	56	7.8	4.3	15.05	1.72
19	B+/B+	128	51	13.1	7.4	8.14	1.48
20	A+/A+	101	51	16.8	9.8	6.86	2.50
21	B+/B+	107	98	4.2	3.7	2.96	0.95
22	O+/A+	109	50	14.0	11.0	13.30	2.64
23	A+/A+	70	36	7.9	4.4	1.10	1.76
24	B+/B+	40	60	1.47	0.7	0.55	2.76
25	O+/A+	108	98	1.80	1.1	1.48	1.14
Mean		82	90	5.38	3.70	5.74	1.81
Median		68	90	4.24	3.00	3.60	1.43
SD		39	35	4.24	2.67	6.00	1.82

\*Conventional thawing techniques were used for Patients 1, 2, and 3, and dextran-albumin thawing techniques for Patients 4 through 25.

†ND denotes not determined.

TABLE 2. PREPARATIVE REGIMENS AND PROPHYLAXIS AGAINST GVHD.

PATIENT NO.	PREPARATIVE REGIMEN*	GVHD PROPHYLAXIS†
1	TBI, melphalan, cyclophosphamide	Methotrexate, cyclosporine, methylprednisolone
2, 6	Busulfan, melphalan	Methotrexate, cyclosporine, methylprednisolone
3	TBI, melphalan	Methotrexate, cyclosporine, methylprednisolone
4	Busulfan, melphalan, cyclophosphamide, ATG	Methotrexate, cyclosporine, methylprednisolone
5, 13	TAI, cyclophosphamide, ATG	Cyclosporine
7, 8, 9, 10, 11, 12	TBI, melphalan, ATG	Methotrexate, cyclosporine, methylprednisolone
14	Busulfan, melphalan, ATG	Cyclosporine
15	TBI, melphalan, ATG	Cyclosporine
16	TBI, melphalan, ATG	Cyclosporine, low-dose methylprednisolone
17	Cyclophosphamide, melphalan, ATG	Cyclosporine, low-dose methylprednisolone
18, 21, 24, 25	TBI, melphalan, ATG	Cyclosporine, high-dose methylprednisolone
19, 22	Busulfan, melphalan, ATG	Cyclosporine, high-dose methylprednisolone
20, 23	Busulfan, cyclophosphamide, ATG	Cyclosporine, high-dose methylprednisolone

\*Patients two years of age or older who had leukemia received 1350 cGy of hyperfractionated total-body irradiation (TBI) on days 8 to 5 before transplantation (days -8 to -5), 60 mg of melphalan per square meter of body-surface area on days -4 to -2, and 30 mg of antithymocyte globulin (ATG; Atgam, Upjohn, Kalamazoo, Mich.) per kilogram on days -3 to -1. In patients under two years of age who had leukemia and Patient 6, who had cardiomyopathy, 16 doses of 40 mg of busulfan per square meter were substituted for TBI. In Patient 17, cyclophosphamide was substituted for TBI. The two patients with Fanconi's anemia were prepared with 15 mg of ATG per kilogram on days -6 to -1, 10 mg of cyclophosphamide per kilogram on days -5 to -2, and 500 cGy of thoracoabdominal irradiation (TAI) on day -1, according to the method of Gluckman et al.<sup>17</sup> Of the three patients with bone marrow failure syndromes or genetic diseases, Patients 20 and 23 were prepared with busulfan on days -9 to -6, 50 mg of cyclophosphamide per kilogram on days -5 to -2, and ATG on days -3 to -1. Patient 4, who had amegakaryocytic thrombocytopenia, had a congenital cardiomyopathy and was given lower-dose cyclophosphamide (1.5 g per square meter) on days -3 and -2, standard-dose busulfan, 90 mg of melphalan per square meter on day -4, and ATG on days -3 to -1.

†Methotrexate, cyclosporine, and methylprednisolone were given as follows: methotrexate, 15 mg per square meter intravenously on day 1 and 10 mg per square meter intravenously on days 3 and 6; intravenous cyclosporine, 5 mg per kilogram on day -2 through day 3, 3 mg per kilogram on days 3 through 15, and 3.75 mg per kilogram on days 16 through 35, followed by oral cyclosporine, 10 mg per kilogram on days 36 through 83, 8 mg per kilogram on days 84 through 97, 6.0 mg per kilogram on days 98 through 119, and 4 mg per kilogram on days 120 through 180; intravenous methylprednisolone, 0.5 mg per kilogram from day 7 through day 14 and 1 mg per kilogram on days 15 through 28, followed by oral methylprednisolone, 0.8 mg per kilogram on days 29 through 42, 0.5 mg per kilogram on days 43 through 56, 0.2 mg per kilogram on days 57 through 119, and 0.1 mg per kilogram on days 120 through 180. The regimen of cyclosporine and low-dose methylprednisolone consisted of 5 mg of cyclosporine per kilogram given intravenously every 12 hours and 30 mg of methylprednisolone per square meter given intravenously on days 5 to 17, with the dose tapered by 10 percent each week thereafter for 8 to 10 weeks, and given intravenously or orally as tolerated by the patient. The regimen of cyclosporine and high-dose methylprednisolone consisted of 5 mg of cyclosporine per kilogram given intravenously every 12 hours and methylprednisolone given intravenously as follows: 10 mg per kilogram on days 5 through 7, 5 mg per kilogram on days 8 through 10, 3 mg per kilogram on days 11 through 13, and 2 mg per kilogram on days 14 through 17; the methylprednisolone dose was tapered by 10 percent each week thereafter and given intravenously or orally as tolerated by the patient.

lin. Busulfan was given instead of total-body irradiation to patients under two years of age who had leukemia. Blood levels were measured after the second of 16 doses to achieve a steady-state concentration of 600 to 900 ng per milliliter, and doses 8 through 16 were modified if necessary. Patients with conditions other than leukemia were treated with other regimens, as described in Table 2. Prophylaxis against GVHD consisted of cyclosporine alone for the two patients with Fanconi's anemia. A combination of methotrexate, cyclosporine, and methylprednisolone was given to the first 11 patients who did not have Fanconi's anemia. Since no case of severe GVHD occurred in these 11 patients, the regimen was reduced to cyclosporine alone in 2 patients and cyclosporine and high-dose methylprednisolone in 10 patients. Methotrexate was given in a dose of 15 mg per square meter of body-surface area on day 1, followed by 10 mg per square meter on days 3 and 6. Leucovorin (15 mg per square meter) was given 24 hours after each dose of methotrexate. The dose of methylprednisolone was 10 mg per kilogram of body weight on days 5 to 7, 5 mg per kilogram on days 8 to 10, 3 mg per kilogram on days 11 and 12, and 2 mg per kilogram on days 15 to 17; thereafter the dose was tapered 10 percent per week. The patients continued to receive a full dose of cyclosporine until

a point between day 180 and day 270 after transplantation, after which the dose was tapered by 10 percent per week.

### Supportive Care

All patients were kept in reverse isolation under high-energy particulate air filtration. A parent roomed with each child. Prophylaxis with trimethoprim-sulfamethoxazole against *Pneumocystis carinii* was initiated before transplantation. Broad-spectrum antibiotic therapy was instituted at the time of the first episode of neutropenic fever and continued until the absolute neutrophil count exceeded 500 per cubic millimeter for two days. All patients received intravenous amphotericin B (0.25 mg per kilogram per day) from day 0 (the day of transplantation). The dose of amphotericin B was increased to 1 mg per kilogram per day if fever persisted for three days after the institution of antibiotic therapy. Patients received transfusions of leukocyte-depleted, irradiated, packed red cells and platelets to maintain platelet counts equal to or greater than 20,000 per cubic millimeter and hematocrit values greater than 27 percent during the first four weeks after transplantation. Filgrastim (10  $\mu$ g per kilogram) was administered daily from day 0

TABLE 3. CHARACTERISTICS OF THE PATIENTS.

PATIENT NO.	AGE AT TRANSPLANTATION yr	SEX	RACE OR ETHNIC GROUP	DIAGNOSIS*	DISEASE STATUS	WEIGHT kg	STATUS BEFORE TRANSPLANTATION
1	3.7	M	White	T-cell ALL	2nd relapse	14.1	History of mediastinal irradiation
2	1.8	M	Black/Hispanic/white	Infant ALL	2nd complete remission	10.1	
3	11.2	M	White	ALL	Relapse	38.5	Never entered complete remission
4	2.6	M	White	AMT		12.9	Transfusion-dependent; congenital cardiomyopathy
5	7.0	F	American Indian/black	FA		17.5	Transfusion-dependent
6	3.5	M	White	ALL	2nd complete remission	19.4	
7	12.0	F	White	T-cell ALL	2nd complete remission	39.1	History of craniospinal irradiation
8	7.8	F	White	KS	ANLL	16.2	Monosomy 7 after filgrastim; aspergillus sinusitis
9	4.1	M	White	ANLL	3rd relapse	16.2	Refractory relapse
10	9.4	M	White	ALL	2nd complete remission	34.0	
11	9.7	M	White	ALL	2nd complete remission	30.5	
12	13.3	M	Black	CML	2nd blast crisis	40.4	
13	23.5	F	American Indian/black	FA	ANLL	44.0	No therapy for ANLL
14	1.2	F	White	Infant ALL	1st complete remission	10.0	
15	5.7	M	White	ALL	2nd complete remission	21.3	
16	11.9	M	White	ALL	2nd complete remission	53.0	
17	2.9	M	White	NB	2nd relapse	12.4	History of mediastinal radiation
18	9.5	M	White/Hispanic	ANLL	1st relapse	24.9	Autologous bone marrow transplantation in 1st complete remission
19	1.3	M	Asian	ANLL	2nd complete remission	12.2	
20	0.8	M	White	LN		7.5	Renal dysfunction
21	11.8	M	Black	ALL	2nd complete remission	32.4	History of hepatosplenic candidiasis
22	1.0	F	White	ANLL	2nd complete remission	8.0	
23	1.1	F	White	CVID-MDS		8.1	Rotavirus infection; polymicrobial sepsis
24	9.3	M	Black	ALL	2nd complete remission	36.4	Pulmonary and hepatosplenic <i>Pneumocystis carinii</i>
25	15.1	M	White	ANLL	2nd relapse	79.0	Autologous bone marrow transplantation in 2nd complete remission

\*ALL denotes acute lymphoblastic leukemia, AMT amegakaryocytic thrombocytopenia, FA Fanconi's anemia, KS Kostmann's syndrome, ANLL acute nonlymphocytic leukemia, CML chronic myelogenous leukemia, NB neuroblastoma, LN Lesch-Nyhan disease, and CVID-MDS common variable immunodeficiency disease-myelodysplastic syndrome.

until the absolute neutrophil count was 10,000 or more per cubic millimeter for three consecutive days or more than 2000 per cubic millimeter for two weeks. Patient 23 had polymicrobial sepsis at the time of preparation for transplantation and was supported with irradiated, filgrastim-mobilized parental granulocytes for the first 10 days after transplantation.

Intravenous immune globulin (Gamimune N, 10 percent, Cutter Biologicals, Elkhart, Ind.) was administered to each patient at a dose of 500 mg per kilogram weekly through day 100 and then once every two weeks or monthly during year 1. Patients who had IgG anti-CMV antibodies before transplantation or who received grafts that were CMV-positive (as determined by culture of the infants' saliva<sup>18</sup>) received ganciclovir through day 100. The presence of IgG anti-CMV antibodies alone in the infant-mother pair was not considered a risk factor for transmission of CMV through the placental-blood graft. Documented CMV infection that occurred after transplantation was treated with therapeutic doses of ganciclovir (5 mg per kilogram per dose, given twice a day) and intravenous immune globulin every other day for three weeks.<sup>19</sup>

#### Study End Points

To evaluate engraftment, the primary end point was the number of days required for the absolute neutrophil count to reach


500 per cubic millimeter. Secondary end points included the demonstration of megakaryocytes in the bone marrow and the number of days needed for the platelet count to remain above 20,000 per cubic millimeter and the hemoglobin level to stay above 10 g per deciliter without transfusion. Chimerism was evaluated by fluorescence in situ hybridization for the X chromosome in sex-mismatched transplants, or DRB1 allele-specific hybridization in cases in which the patient and donor differed at HLA-DR. These tests were performed 28 to 35, 100, 180, and 360 days after transplantation. Complete chimerism was inferred when all cells in the marrow and peripheral blood of the patient were of donor origin, whereas mixed chimerism was defined as the simultaneous presence of both donor and host cells. Primary graft failure was defined as a failure to reach a white-cell count of 500 per cubic millimeter or an absolute neutrophil count of 200 per cubic millimeter within 30 days of transplantation or a continued need for platelet transfusions for more than 100 days after transplantation in the absence of a frank leukemic relapse. Immunologic studies included assays for lymphocyte proliferation in response to plant mitogens (phytohemagglutinin, concanavalin A, and pokeweed mitogen) and tetanus toxoid and *Candida albicans* antigens; counting of the lymphocyte subgroups by fluorescence-activated cell sorting; and an assay of natural-killer-cell function by the measurement of lysis of the K562 cell line.

TABLE 4. HLA TYPING OF DONORS AND RECIPIENTS.\*

PATIENT NO. AND BLOOD UNIT	CLASS I (SEROLOGIC TYPING)		CLASS II (LOW-RESOLUTION TYPING)	CLASS II (HIGH-RESOLUTION TYPING)	NO. OF MISMATCHED ANTIGENS	
	A	B	DR	DRB1	LOW RESOLUTION	HIGH RESOLUTION
	antigens		gene groups	genes		
Patient 1	2,29	<b>44,60</b>	4,7	<b>0401,0701</b>		
Blood unit	2,29	<b>44,50</b>	4,7	<b>0407,0701</b>	1	2
Patient 2	2,25	50,57	4,7	<b>04xx†,0701</b>		
Blood unit	2,26	50,57	<b>blank,7</b>	0701,0701	1	2
Patient 3	2,28	44,51	<b>8,11</b>	<b>0801,1101</b>		
Blood unit	2,28	44,51	7,11	<b>0701,1101</b>	1	1
Patient 4	2,blank	<b>blank,44</b>	12,13	1201,1301		
Blood unit	2,blank	<b>41,44</b>	12,13	1201,1302	1	2
Patient 5	11,26	<b>35,70</b>	2,4	<b>1503,0401</b>		
Blood unit	11,26	<b>38,70</b>	2,4	<b>1501,0402</b>	1	3
Patient 6	2,blank	<b>57,62</b>	2,7	1501,0701		
Blood unit	2,blank	<b>57,44</b>	2,7	<b>1601,0701</b>	1	2
Patient 7	24,28	<b>39,60</b>	4,blank	<b>0403,0407</b>		
Blood unit	24,28	<b>39,61</b>	4,blank	<b>0411,0407</b>	1	2
Patient 8	2,32	<b>7,blank</b>	1,2	<b>0101,1501</b>		
Blood unit	2,1	<b>7,blank</b>	1,2	<b>0103,1501</b>	1	2
Patient 9	2,3	7,18	7,11	0701,1101		
Blood unit	2,3	<b>7,50</b>	7,11	0701,1101	1	1
Patient 10	1,30	13,37	7,11	0701,1103		
Blood unit	1,30	13,37	7,11	0701,1103	0	0
Patient 11	11,31	<b>35,40</b>	4,blank	<b>0404,blank</b>		
Blood unit	11,31	<b>35,58</b>	4,blank	<b>0402,blank</b>	1	2
Patient 12	2,3	<b>7,58</b>	<b>13,blank</b>	<b>1301,1303</b>		
Blood unit	2,3	<b>7,39</b>	<b>2,8</b>	<b>0801,1501</b>	3	3
Patient 13	28,30	<b>42,52</b>	3,8	0302,0804		
Blood unit	28,30	7,42	3,8	0302,0804	1	1
Patient 14	2,blank	27,50	2,7	1501,0701		
Blood unit	2,blank	27,50	<b>7,blank</b>	<b>0701,blank</b>	1	1
Patient 15	11,24	14,51	1,11	0102,1104		
Blood unit	11,24	14,51	1,13	0102,1301	1	1
Patient 16	1,2	<b>8,60</b>	3,4	0301,0404		
Blood unit	1,2	<b>8,44</b>	3,4	0301,0401	1	2
Patient 17	3,32	8,blank	1,2	0101,1501		
Blood unit	3,2	8,blank	1,2	0101,1501	1	1
Patient 18	2,blank	<b>35,62</b>	1,14	0101,1402		
Blood unit	2,blank	<b>27,62</b>	1,4	0101,0401	2	2
Patient 19	2,blank	<b>15,46</b>	2,12	1201,1501		
Blood unit	2,11	<b>15,13</b>	2,12	1201,1502	2	3
Patient 20	24,33	14,62	1,4	0102,0404		
Blood unit	24,3	14,62	1,4	0102,0404	1	1
Patient 21	3,31	7,14	1,4	0102,0404		
Blood unit	3,33	7,14	1,4	0102,0404	1	2
Patient 22	2,29	44,62	4,7	0401,0701		
Blood unit	1,29	44,62	4,7	0401,0701	1	1
Patient 23	2,30	<b>13,52</b>	2,11	<b>1103,1501</b>		
Blood unit	2,30	<b>35,52</b>	2,11	<b>1104,1502</b>	1	3
Patient 24	3,30	7,58	2,11	1503,1101		
Blood unit	3,28	<b>49,58</b>	2,11	1503,1101	2	2
Patient 25	1,3	7,8	3,7	0301,0701		
Blood unit	1,3	<b>7,14</b>	3,7	0301,0701	1	1
	<b>NO. OF MISMATCHED ANTIGENS</b>	<b>NO. OF TRANSPLANTS MISMATCHED BY LOW-RESOLUTION TYPING</b>	<b>NO. OF TRANSPLANTS MISMATCHED BY HIGH-RESOLUTION TYPING</b>			
	0	1	1			
	1	20	9			
	2	3	11			
	3	1	4			

\*Mismatches are indicated by boldface type.

†Novel DR4 genotype.



**Figure 1.** Relation of Neutrophil Recovery to Thawing Method. Values shown are means  $\pm$  SD.

### GVHD

GVHD was scored according to standard criteria.<sup>9</sup> During the first month after transplantation, all the patients were evaluated daily. After discharge from the hospital, patients were seen twice weekly during the second month after transplantation, weekly during the third through sixth months, and then quarterly. GVHD was documented histopathologically. DNA probes for maternal HLA genes were used to seek grafted cells derived from the donor's mother in all biopsy samples. GVHD (of grade II or higher) was treated with a high dose of methylprednisolone for three to seven days, after which the dose was tapered.

### Statistical Analysis

The probability of event-free survival was calculated by Kaplan-Meier analysis.<sup>20</sup> Data on patients were censored at the first adverse event, such as death due to treatment-related toxicity, graft failure, relapse, or death from other causes. Correlations between the numbers of nucleated, hematopoietic progenitor, and CD34+ cells infused and the length of time to engraftment were expressed as the linear correlation coefficient ( $R^2$ ) and evaluated by the t-test with SPSS for Windows software (SPSS, Chicago).

## RESULTS

### Patient Characteristics

Twenty-five patients received transplants from August 1993 through November 1995 (Table 3). Searches for matched, unrelated bone marrow donors failed to identify such a donor for 17 of 22 patients. The median weight of the patients was 19.4 kg (range, 7.5 to 79.0) and the median age was 7.0 years (range, 0.8 to 23.5). Nineteen patients had malignant diseases and four had nonmalignant conditions; in two patients (Patients 8 and 23), a primary nonmalignant condition had converted to malignant disease. In nine patients, prior infection with CMV was evidenced by IgG anti-CMV antibody titers of 1:8 or higher. Two patients with acute non-lymphocytic leukemia underwent transplantation during a relapse after they had undergone autologous bone marrow transplantation.

### Selection of Units of Placental Blood for Transplantation

Units of placental blood were initially matched to the patient's HLA phenotype by serologic typing for

class I HLA antigens and low-resolution DNA typing for class II HLA alleles. By these means, 1 of the 25 units we transplanted was matched for six of six HLA antigens, 20 were matched for five of six, 3 for four of six, and 1 for three of six. Subsequently, all the patients and units of placental blood were typed for HLA-DRB1 by high-resolution DNA hybridization with group-specific polymerase-chain-reaction primers and allele-specific oligonucleotide probes. This second genetic analysis revealed the following distribution of HLA matches: 1 donor-recipient pair at six loci, 9 pairs at five loci, 11 pairs at four loci, and 4 pairs at three loci (Table 4). When a matched unit was available, the median time to its identification was 3 days, and the median time to transplantation 102 days. When a continued search was required to find a suitable unit, the average time to identification was 18 days (range, 0 to 128), and the time to transplantation 115 days (range, 12 to 291).

### Engraftment

There was evidence of myeloid engraftment in 23 of the 25 patients. In one patient (Patient 3), the infused cells failed to engraft, but spontaneous reconstitution with autologous cells occurred 65 days after transplantation. Two patients (Patients 9 and 12) had persistent leukemia (one of them had evidence of myeloid engraftment), and the absolute neutrophil count never rose above 500 per cubic millimeter. In the remaining 22 patients, the absolute neutrophil count reached 500 per cubic millimeter in a median of 22 days (range, 14 to 37). Platelet transfusions became unnecessary in a median of 56 days (range, 35 to 89) in 16 patients who could be evaluated. Platelet counts of 50,000 and 100,000 per cubic millimeter were reached by a median of 82 and 115 days, respectively. Red-cell transfusions could be stopped after a median of 55 days (range, 32 to 90). Seven patients died of relapse (Patient 9), regimen-related toxicity (Patients 7 and 17), or infection (Patients 8, 11, 19, and 21) while still dependent on platelet or red-cell transfusions. All surviving patients were complete chimeras as of the most recent follow-up in May 1996.

### Cell Dose and Speed of Engraftment

The number of nucleated cells infused per kilogram of the patient's body weight correlated with the rate of myeloid engraftment ( $P=0.002$ ; data not shown). There was a trend for the time to myeloid or platelet engraftment to increase with the dose of clonogenic precursors or CD34+ cells, but the correlations were not statistically significant. After observing delayed recovery of neutrophils in the first three patients, we began to use a new thawing technique that increases cell viability in vitro.<sup>16</sup> In patients infused with these "washed" units of placental blood, myeloid engraftment was acceler-

TABLE 5. IMMUNOLOGIC RECONSTITUTION.

PATIENT No.	POST-TRANSPLANTATION DAY	MITOGEN*			ABSOLUTE LYMPHOCYTE COUNT	CD4:CD8 RATIO
		PHA	CON A	PWM		
		counts per minute			cells/mm <sup>3</sup>	
1	60	141,124	62,712	53,226	1,408	0.33
2	95	63,561	53,968	26,011	2,176	2.25
	192	59,991	42,024	26,817	2,279	0.94
3	380	245,573	279,177	109,366	2,262	1.46
	820	153,606	280,548	85,583	5,568	1.93
	118	128,537	139,027	81,733	495	0.80
4	245	105,815	122,060	40,860	690	3.99
	361	163,680	417,051	276,527	986	2.07
	418	298,166	239,396	33,580	1,350	2.04
5	84	92,050	90,444	2,085	1,463	0.33
	176	101,962	53,207	22,033	2,272	0.60
	260	66,128	35,254	2,291	9,570	1.05
	358	111,070	59,867	3,567	11,900	1.42
	512	276,131	324,249	59,229	4,620	0.78
6	95	90,539	81,933	17,245	469	0.37
	158	82,141	114,941	46,765	1,504	0.67
10	71	58,656	76,793	8,705	792	0.54
	204	166,290	116,710	47,095	NA	0.64
	428	284,755	237,501	143,944	2,624	1.92
13	95	87,094	79,724	14,797	266	0.58
14	60	22,617	39,788	1,614	4,020	0.11
	103	47,114	52,213	30,078	3,000	2.47
	299	307,475	208,247	21,369	4,092	0.78
15	54	248,565	254,876	13,328	510	2.33
	98	146,864	90,537	10,900	900	1.42
16	190	192,657	197,215	52,592	736	0.87
20	187	264,935	189,613	156,324	2,720	1.93
22	95	61,829	58,982	4,280	940	2.31
23	53	146,650	74,027	137,584	648	1.51

\*PHA denotes phytohemagglutinin, Con A concanavalin A, and PWM pokeweed mitogen. The respective mean ( $\pm$ SD) normal values in children from the Duke University Medical Center Clinical Pediatric Immunology Laboratory are  $240,684 \pm 125,036$ ,  $214,779 \pm 112,333$ , and  $157,469 \pm 60,364$ . NA denotes not available.

ated (Fig. 1), but platelet and red-cell engraftment was not affected.

#### GVHD

Twenty-one of the 25 patients could be evaluated for GVHD. Four patients did not have GVHD; eight patients had grade I GVHD, involving only the skin; and seven patients had grade II GVHD, involving the skin and gut. In two patients (Patients 13 and 16, who were recipients of grafts that were mismatched at HLA-B and at HLA-B and HLA-DRB1, respectively), grade III GVHD, involving the skin and gut, developed. No patient had acute grade IV GVHD. In two patients (Patients 5 and 16), chronic GVHD limited to the liver or skin developed 19 and 7 months after transplantation. There was no correlation between the incidence or extent of GVHD and the degree of HLA mismatching (data not shown). Maternal cells from the donor unit were not found in the tissue-biopsy specimens from any patient with GVHD. All patients with GVHD

responded to treatment with high doses of methylprednisolone, and none required second-line therapy for steroid-refractory GVHD.

#### Immunologic Reconstitution

In vitro responses to T-cell and B-cell mitogens were detectable in 13 patients with engraftment who were studied within three months of transplantation (Table 5). The absolute lymphocyte counts were greater than 500 per cubic millimeter, but the CD4:CD8 ratios were inverted in all patients studied for the first six months after transplantation. Natural-killer-cell function was normal in six patients tested two to three months after transplantation.

#### Survival

As of June 1996, 12 of the 25 patients had survived event-free for 7 to 32 months after transplantation (Table 6), for an event-free-survival rate of 48 percent. Seven of the 19 patients undergoing transplantation for malignant conditions and 5 of the

6 with nonmalignant conditions had survived event-free with Karnofsky scores above 90, with a median follow-up of 12½ months (range, 7 to 32). Only one of the eight patients who underwent transplantation while in remission and survived for more than 100 days had had a leukemic relapse. Thirteen of the 25 patients had died of infection, relapse, or toxic effects of treatment.

**DISCUSSION**

In this study we found that hematopoietic stem cells in placental blood from unrelated donors engrafted and reconstituted hematopoiesis in more than half of a group of high-risk patients, nearly all of them children, with malignant and nonmalignant conditions. There was a complete match between the donor and the recipient for all class I HLA antigens and class II HLA alleles in only one case. Despite the HLA incompatibility, GVHD was mild (less

than grade III) in all but two recipients. The substitution of melphalan for cyclophosphamide in patients with malignant conditions allowed adequate engraftment.<sup>22-29</sup> Complete donor chimerism was achieved without total-body irradiation in nine patients who were prepared with chemotherapy alone. This experience is relevant for transplantation in younger children, who are at especially high risk for treatment-related neurotoxicity.<sup>30</sup> The recovery of platelets and red cells was later in all recipients of placental-blood grafts than in recipients of marrow grafts, perhaps because placental blood contains a higher proportion of immature hematopoietic progenitor cells than does bone marrow.<sup>31-38</sup>

Engraftment occurred in patients who received as few as 6 million nucleated, HLA-disparate, placental-blood cells per kilogram, but we do not know whether this result in children can be applied to adults.<sup>31-38</sup> We found a relation between the speed of

**TABLE 6.** ENGRAFTMENT, GVHD, AND SURVIVAL IN PATIENTS WHO RECEIVED HLA-MISMATCHED PLACENTAL BLOOD.

PATIENT NO.	DATE OF TRANSPLANTATION	WEIGHT kg	CELL DOSE ×10 <sup>-7</sup> /kg	No. OF MISMATCHES		TBI*	ENGRAFTMENT	GVHD GRADE†	EFS‡	CAUSE OF DEATH OR EVENT
				LOW RESOLUTION	HIGH RESOLUTION					
1	8/24/93	14.1	2.7	1	2	+	Yes	0	61	Interstitial pneumonia
2	9/13/93	10.1	3.7	1	2	-	Yes	II	>998	
3	3/9/94	38.5	0.7	1	1	+	No	NE	30	Graft failure
4	8/19/94	12.9	6.8	1	2	-	Yes	II	>658	
5	9/27/94	17.5	2.1	1	3	+§	Yes	II	>619	
6	10/27/94	19.4	4.8	1	2	-	Yes	II	180	Relapse
7	11/9/94	39.1	1.3	1	2	+	Yes	II	46	Pulmonary alveolar hemorrhage
8	11/15/94	16.2	2.1	1	2	+	Yes	NE	15	Aspergillus
9	12/14/94	16.1	2.4	1	1	+	Yes	NE	41	Relapse
10	1/5/95	34.0	1.5	0	0	+	Yes	0	>519	
11¶	1/17/95	30.5	2.0	1	2	+	Yes	0	55	Toxoplasmosis
12	2/1/95	40.4	1.6	3	3	+	No	NE	30	Graft failure, interstitial pneumonia
13	2/24/95	43.0	3.7	1	1	+‡	Yes	III	>469	
14	4/28/95	10.0	5.4	1	1	-	Yes	I	>406	
15	6/14/95	21.3	3.0	1	1	+	Yes	I	>359	
16	7/5/95	53.0	3.8	1	2	+	Yes	III	>338	
17	8/7/95	12.7	2.4	1	1	-	Yes	0	35	Interstitial pneumonia
18	8/9/95	24.9	4.3	2	2	+	Yes	I	>303	
19	8/22/95	12.2	7.4	2	3	-	Yes	II	56	CMV
20	8/23/95	7.5	9.8	1	1	-	Yes	I	>289	
21	9/6/95	32.4	3.7	1	2	+	Yes	II	77	Aspergillus
22	10/13/95	8.0	11.0	1	1	-	Yes	I	>238	
23	10/26/95	8.1	4.4	1	3	-	Yes	II	>225	
24	11/8/95	36.4	0.7	2	2	+	Yes	II	162	Pulmonary failure
25	11/8/95	79.0	1.1	1	1	+	Yes	I	103	Adenovirus

\*A plus sign indicates that the patient underwent total-body irradiation (TBI), and a minus sign that the patient did not.

†NE denotes not evaluated.

§The patient received thoracoabdominal irradiation in place of TBI.

||Patient 12 had persistent chronic myelogenous leukemia and graft failure and received a second placental-blood graft.

‡Event-free survival (EFS) is given through June 7, 1996.

¶The placental-blood graft was negative for toxoplasmosis.<sup>21</sup>

myeloid recovery and the dose of infused cells, but the International Cord Blood Transplant Registry found no such correlation in recipients of placental blood from related donors.<sup>11</sup> This difference may be due to the uniformity of results a single center can obtain, or to the use of filgrastim in our patients. Although the number of CD34+ cells (presumably hematopoietic progenitor cells) in the graft correlates with myeloid and platelet engraftment in recipients of filgrastim-mobilized peripheral-blood progenitor cells, we did not find any such correlation.

All but one of the pairs of donors and recipients in this series differed by one to three HLA antigens. Nevertheless, primary graft failure occurred only in patients who underwent transplantation during leukemic relapse, and severe acute or chronic GVHD was not observed. By contrast, in 462 patients undergoing bone marrow transplantation from HLA-matched, unrelated donors, the probability of grade III or IV acute GVHD was 47 percent overall and 30 percent for patients less than 18 years of age; for chronic GVHD, the probability was 55 percent.<sup>2</sup> Contamination of placental blood by maternal cells is a potential cause of serious GVHD,<sup>39-43</sup> but our results do not support this possibility. The observation that placental T cells mount less of a graft-versus-host response than bone marrow from unrelated donors in HLA-mismatched recipients has yet to be explained, but it parallels observations made in vitro and in animal studies.<sup>44</sup> Another unsolved problem is whether the lower incidence of GVHD indicates a decrease in graft-versus-leukemia activity and thus an increase in the risk of leukemic relapse. Our data are too premature to answer this question.

Banked placental blood has potential advantages over bone marrow from adults. Placental blood is less likely than adult bone marrow to contain viruses; the storage of placental blood reduces procurement time to one or two weeks, substantially less than the four to six months typically needed to find an unrelated bone marrow donor; and with placental blood the time required to prepare the patient for transplantation is usually short.

A major disadvantage of placental-blood transplantation is that only one unit is available for each transplantation procedure. Ex vivo expansion of stem cells and progenitor cells might circumvent this problem. A second possible disadvantage is the unwitting transmission of a genetic disease affecting hematopoietic cells. Placental blood can be tested for common hematologic diseases, such as hemoglobinopathies, before banking or transplantation, and the family history can reveal the possibility of an inherited disorder. Nevertheless, extremely rare genetic diseases are unlikely to be revealed by the family history, and laboratory testing for them may be impractical or impossible.

In summary, we have demonstrated that partially

mismatched placental blood from unrelated donors can provide an alternative source of stem cells for hematopoietic reconstitution. The use of placental blood that differed from the recipient's by one to three HLA alleles resulted in 100 percent donor chimerism, generally treatable GVHD, and immune reconstitution.

Supported in part by a grant (1R18-HL48031) from the National Heart, Lung, and Blood Institute and by the Leukemia Society of America.

*We are indebted to the nursing staff of the Pediatric Bone Marrow Transplant Program at Duke University Medical Center for caring for these patients with diligence and compassion; to Alice Stewart and the staff of the Pediatric Bone Marrow Transplant Laboratory for characterizing and processing the placental-blood units; and to Jill Beindick and Connie Stephens for data collection and coordinating the searches, sample procurement, and engraftment studies; to the referring physicians (Tanya Trippett, Charles Daeschner, Kenneth Starling, Deborah Scott, Felicia Little, Eric Werner, Roger Berkow, Bob Ettinger, Raj Malik, Barry Golemb, Allen Chauvenet, Paul Martin, Ming Yang, Virgil Rose, Mark Mogul, Gary Jones, Stuart Gold, Frank Keller, Kim Ritchie, Richard Schiff, Terry Harville, and Rebecca Buckley) for their willingness to refer their patients and for their enthusiasm and compliance in supplying follow-up data; and to the staff of the Department of Obstetrics and Gynecology and the Department of Neonatology at Mount Sinai Medical Center, New York, for their support in the collection of blood units and samples from babies and mothers.*

## REFERENCES

1. Gluckman E. Bone marrow transplantation in Fanconi's anemia. *Stem Cells* 1993;11:Suppl 2:180-3.
2. Kernan NA, Bartsch G, Ash RC, et al. Analysis of 462 transplantations from unrelated donors facilitated by the National Marrow Donor Program. *N Engl J Med* 1993;328:593-602.
3. Storb R, Thomas ED, Buckner CD, et al. Marrow transplantation for aplastic anemia. *Semin Hematol* 1984;21:27-35.
4. Champlin RE, Ho WG, Nimer SD, et al. Bone marrow transplantation for severe aplastic anemia: effect of a preparative regimen of cyclophosphamide-low-dose total-lymphoid irradiation and posttransplant cyclosporine-methotrexate therapy. *Transplantation* 1990;49:720-4.
5. McGlave PB, Haake R, Miller W, Kim T, Kersey J, Ramsay NK. Therapy of severe aplastic anemia in young adults and children with allogeneic bone marrow transplantation. *Blood* 1987;70:1325-30.
6. Storb R, Sanders JE, Pepe M, et al. Graft-versus-host disease prophylaxis with methotrexate/cyclosporine in children with severe aplastic anemia treated with cyclophosphamide and HLA-identical marrow grafts. *Blood* 1991;78:1144-5.
7. Storb R, Pepe M, Anasetti C, et al. What role for prednisone in prevention of acute graft-versus-host disease in patients undergoing marrow transplants? *Blood* 1990;76:1037-45.
8. Chao NJ, Schmidt GM, Niland JC, et al. Cyclosporine, methotrexate, and prednisone compared with cyclosporine and prednisone for prophylaxis of acute graft-versus-host disease. *N Engl J Med* 1993;329:1225-30.
9. Przepiorka D, Weisdorf D, Martin P, et al. 1994 Consensus Conference on Acute GVHD Grading. *Bone Marrow Transplant* 1995;15:825-8.
10. Howard MR, Gore SM, Hows JM, Downie TR, Bradley BA. A prospective study of factors determining the outcome of unrelated marrow donor searches: report from The International Marrow Unrelated Search and Transplant Study Working Group on behalf of collaborating centres. *Bone Marrow Transplant* 1994;13:389-96.
11. Wagner JE, Kernan NA, Steinbuch M, Broxmeyer HE, Gluckman E. Allogeneic sibling umbilical-cord-blood transplantation in children with malignant and non-malignant disease. *Lancet* 1995;346:214-9.
12. Gluckman E, Broxmeyer HE, Auerbach AD, et al. Hematopoietic reconstitution in a patient with Fanconi's anemia by means of umbilical-cord blood from a HLA-identical sibling. *N Engl J Med* 1989;321:1174-8.
13. Kurtzberg J, Graham M, Casey J, Olson J, Stevens CE, Rubinstein P. The use of umbilical cord blood in mismatched related and unrelated hematopoietic stem cell transplantation. *Blood Cells* 1994;20:275-84.
14. Vilmer E, Sterkers G, Rahimy C, et al. HLA-mismatched cord blood

- transplantation in a patient with advanced leukemia. *Bone Marrow Transplant* 1991;7:Suppl 2:125.
15. Rubinstein P, Rosenfield RE, Adamson JW, Stevens CE. Stored placental blood for unrelated bone marrow reconstitution. *Blood* 1993;81:1679-90.
  16. Rubinstein P, Dobrila L, Rosenfield RE, et al. Processing and cryopreservation of placental/umbilical cord blood for unrelated bone marrow reconstitution. *Proc Natl Acad Sci U S A* 1995;92:10119-22.
  17. Gluckman E, Socie G, Devergie A, Bourdeau-Esperou H, Traineau R, Cosset JM. Bone marrow transplantation in 107 patients with severe aplastic anemia using cyclophosphamide and thoraco-abdominal irradiation for conditioning: long-term follow-up. *Blood* 1991;78:2451-5.
  18. Balcarek KB, Warren W, Smith RJ, Lyon MD, Pass RF. Neonatal screening for congenital cytomegalovirus infection by detection of virus in saliva. *J Infect Dis* 1993;167:1433-6.
  19. Emanuel D, Cunningham I, Jules-Elysee K, et al. Cytomegalovirus pneumonia after bone marrow transplantation successfully treated with the combination of ganciclovir and high-dose intravenous immune globulin. *Ann Intern Med* 1988;109:777-82.
  20. Kaplan EL, Meier P. Nonparametric estimation from incomplete observations. *J Am Stat Assoc* 1958;53:457-81.
  21. Peacock JE Jr, Greven CM, Cruz JM, Hurd DD. Reactivation toxoplasmic retinochoroiditis in patients undergoing bone marrow transplantation: is there a role for chemoprophylaxis? *Bone Marrow Transplant* 1995;15:983-7.
  22. Samuels BL, Bitran JD. High-dose intravenous melphalan: a review. *J Clin Oncol* 1995;13:1786-99.
  23. Powles RL, Milliken S, Helenglass G, et al. The use of melphalan in conjunction with total body irradiation as treatment for acute leukaemia. *Transplant Proc* 1989;21:2955-7.
  24. Carey PJ, Proctor SJ, Taylor P, Hamilton PJ. Autologous bone marrow transplantation for high-grade lymphoid malignancy using melphalan/irradiation conditioning without marrow purging or cryopreservation. *Blood* 1991;77:1593-8.
  25. Gandola L, Lombardi F, Siena S, et al. Total body irradiation and high-dose melphalan with bone marrow transplantation at Istituto Nazionale Tumori, Milan, Italy. *Radiother Oncol* 1990;18:Suppl 1:105-9.
  26. Schroeder H, Pinkerton CR, Powles RL, et al. High dose melphalan and total body irradiation with autologous marrow rescue in childhood acute lymphoblastic leukaemia after relapse. *Bone Marrow Transplant* 1991;7:11-5.
  27. Keating A, Crump M. High dose etoposide melphalan, total body irradiation and ABMT for acute myeloid leukemia in first remission. *Leukemia* 1992;6:Suppl 4:90-1.
  28. Phillips GL, Shepherd JD, Barnett MJ, et al. Busulfan, cyclophosphamide, and melphalan conditioning for autologous bone marrow transplantation in hematologic malignancy. *J Clin Oncol* 1991;9:1880-8.
  29. Russell S, Vowels M. Busulfan, cyclophosphamide, and melphalan as conditioning therapy in allogeneic bone marrow transplants for acute lymphoblastic leukemia. *Transplant Proc* 1992;24:183.
  30. Wingard JR, Plotnick LP, Freemer CS, et al. Growth in children after bone marrow transplantation: busulfan plus cyclophosphamide versus cyclophosphamide plus total body irradiation. *Blood* 1992;79:1068-73.
  31. Broxmeyer HE. Self-renewal and migration of stem cells during embryonic and fetal hematopoiesis: important, but poorly understood events. *Blood Cells* 1991;17:282-6.
  32. Broxmeyer HE, Douglas GW, Hangoc G, et al. Human umbilical cord blood as a potential source of transplantable hematopoietic stem/progenitor cells. *Proc Natl Acad Sci U S A* 1989;86:3828-32.
  33. Broxmeyer HE, Hangoc G, Cooper S, et al. Growth characteristics and expansion of human umbilical cord blood and estimation of its potential for transplantation in adults. *Proc Natl Acad Sci U S A* 1992;89:4109-13.
  34. Lu L, Xiao M, Shen RN, Grigsby S, Broxmeyer HE. Enrichment, characterization, and responsiveness of single primitive CD34 human umbilical cord blood hematopoietic progenitors with high proliferative and replating potential. *Blood* 1993;81:41-8.
  35. Hows JM, Bradley BA, Marsh JCW, et al. Growth of human umbilical cord blood in longterm haemopoietic cultures. *Lancet* 1992;340:73-6.
  36. Hirao A, Kawano Y, Takaue Y, et al. Engraftment potential of peripheral and cord blood stem cells evaluated by a long-term culture system. *Exp Hematol* 1994;22:521-6.
  37. Migliaccio G, Migliaccio AR, Druzin ML, Giardina PJ, Zsebo KM, Adamson JW. Long-term generation of colony-forming cells in liquid culture of CD34+ cord blood cells in the presence of recombinant human stem cell factor. *Blood* 1992;79:2620-7.
  38. Gabutti V, Timeus F, Ramenghi U, et al. Human cord blood progenitors: kinetics, regulation and their use for hemopoietic reconstitution. *Bone Marrow Transplant* 1993;12:Suppl 1:84-6.
  39. Wang XH, Zipursky A. Maternal erythrocytes in the fetal circulation: the immunocytochemical identification of minor populations of erythrocytes. *Am J Clin Pathol* 1987;88:346-8.
  40. Cohen F, Zuelzer WW. The transplacental passage of maternal erythrocytes into the fetus. *Am J Obstet Gynecol* 1965;93:566-9.
  41. O'Reilly RJ, Patterson JH, Bach FH, et al. Chimerism detected by HL-A typing. *Transplantation* 1973;15:505-7.
  42. Pollack MS, Kapoor N, Sorell M, et al. DR-positive maternal engrafted T cells in a severe combined immunodeficiency patient without graft-versus-host disease. *Transplantation* 1980;30:331-4.
  43. Pollack MS, Kirkpatrick D, Kapoor N, Dupont B, O'Reilly RJ. Identification by HLA typing of intrauterine-derived maternal T cells in four patients with severe combined immunodeficiency. *N Engl J Med* 1982;307:662-6.
  44. Risdon G, Gaddy J, Stehman FB, Broxmeyer HE. Proliferative and cytotoxic responses of human cord blood T lymphocytes following allogeneic stimulation. *Cell Immunol* 1994;154:14-24.

## CORRECTION

## Placental-Blood Transplantation

*To the Editor:* Substantial direct and indirect harm may occur during the removal of cord blood from infants. Large amounts of blood are needed to obtain sufficient numbers of stem cells for transplantation. The procedure for removing cord blood can be manipulated to obtain more cord blood, leaving infants anemic. Before stem cells are transplanted, they must be screened for genetic and infectious diseases, including infection with the human immunodeficiency virus (HIV). Although ethical issues concerning the need to inform parents of diseases that have been detected in their infants have been raised, they have not been resolved. Indeed, the recent issuance of a European patent on cord-blood stem cells and the existence of three or more commercial companies to harvest stem cells have increased the demand to examine ethical issues.<sup>1</sup> We need only remember the recent criticism of anonymous testing of infants for HIV because of the failure to inform parents of the diagnosis and offer treatment.<sup>2</sup>

Undoubtedly, repositories of cord-blood stem cells will be widely established. As Silberstein and Jefferies note in their editorial (July 18 issue),<sup>3</sup> the potential benefits to adults, society, science, and commercial interests are substantial and include the use of allogeneic stem-cell therapy for malignant conditions and infectious diseases, the discovery of new genetic disorders, the issuing of patents for newly discovered genes, and profit from therapeutic uses.<sup>4</sup> However, there is a major gap in drug development between children and adults. Children often do not benefit from drugs developed for adults, and there are now proposals to take cells from infants largely for the benefit of adults.

Infants must be the first to benefit in some way from the use of their own stem cells. Repositories must be called by their scientifically correct name, "infant-stem-cell repositories," not placental repositories. A permanent record of diseases that are diagnosed in the infants should be maintained. Parents should be informed of the diagnoses. Therapy should be offered for treatable diseases. The commercialization of stem cells from infants for therapy should include financial benefit for the infants. Royalties from each stem-cell donation should be placed in a national trust fund to be used for research and the treatment of children with serious and life-threatening diseases. The rights of infants as donors cannot be ignored.

Arthur J. Ammann, M.D.  
University of California, San Francisco  
San Francisco, CA 94143

## References

1. Butler D. US company comes under fire over patent on umbilical cord cells. *Nature* 1996;382:99-99.

2. Sacrificing babies on the altar of privacy. *Wall Street Journal*. August 3, 1995.
3. Silberstein LE, Jefferies LC. Placental-blood banking – a new frontier in transfusion medicine. *N Engl J Med* 1996;335:199-201.
4. Sugarman J, Reisner EG, Kurtzberg J. Ethical aspects of banking placental blood for transplantation. *JAMA* 1995;274:1783-1785.

*To the Editor:* The article by Kurtzberg et al. (July 18 issue)<sup>1</sup> describes 25 children who received placental blood from unrelated donors. The median weight of these patients was 19.4 kg (range, 7.5 to 79.0), and they received a mean of 1.81 million CD34+ cells per kilogram of body weight (median, 1.43 million; SD, 1.82 million). This is in sharp contrast to observations by us<sup>2,3</sup> and others<sup>4</sup> that placental-blood units collected for allogeneic bone marrow reconstitution contain 500,000 to 5 million CD34+ cells, a population that includes hematopoietic progenitor cells. We believe that this point should be clarified, particularly because the calculated number of CD34+ cells per kilogram is increasingly used to evaluate the engraftment potential of circulating hematopoietic progenitor cells collected for autologous or allogeneic transplantation.

Francesco Bertolini, M.D., Ph.D.  
Gioacchino Robustelli della Cuna, M.D.  
*I.R.C.C.S. Maugeri Foundation*  
27100 Pavia, Italy

## References

1. Kurtzberg J, Laughlin M, Graham ML, et al. Placental blood as a source of hematopoietic stem cells for transplantation into unrelated recipients. *N Engl J Med* 1996;335:157-166.
2. Bertolini F, Lazzari L, Lavri E, et al. Comparative study of different procedures for the collection and banking of umbilical cord blood. *J Hematother* 1995;4:29-36.
3. Bertolini F, Battaglia M, Zibera C, et al. A new method for placental/cord blood processing in the collection bag. I. Analysis of factors involved in red blood cell removal. *Bone Marrow Transplant* 1996;18:783-786.
4. Kögler G, Callejas J, Hakenberg P, et al. Hematopoietic transplant potential of unrelated cord blood: critical issues. *J Hematother* 1996;5:105-116.

*To the Editor:* We agree with Kurtzberg et al. that in recipients of cord blood the recovery of T lymphocytes is usually rapid. However, the prompt recovery of lymphocyte function in vitro does not necessarily correlate with effective in vivo immunity, because a significant number of children treated by Kurtzberg et al. died of opportunistic infections. We suggest that the results of in vitro studies of lymphocyte function require cautious interpretation.

In our four recipients of cord blood we observed an impressive increase in the percentage (up to 50 percent of the lymphocyte population) and absolute number (up to 5000 per cubic millimeter) of CD19+ B cells, which was apparently not related to defective control of Epstein-Barr virus infection, from about two to nine months after cord-blood transplantation. Most of these lymphocytes also expressed CD20 and surface IgM. Moreover, serum levels of IgM and IgA were those expected in children less than one year of age. Recipients of cord blood might repeat the neonate's program of B-cell ontogeny, with antigenic challenge favoring an expansion in the population of B lymphocytes.<sup>1</sup> In this regard, it is known that memory B cells reside in adult bone marrow, but that cord blood contains only naive B lymphocytes.

Franco Locatelli, M.D.

Rita Maccario, Ph.D.

Marco Zecca, M.D.

*I.R.C.C.S. Policlinico San Matteo*

*27100 Pavia, Italy*

## References

1. Vitetta ES, Berton MT, Burger C, Kepron M, Lee WT, Yin XM. Memory B and T cells. *Annu Rev Immunol* 1991;9:193-217.

*To the Editor:* Kurtzberg et al. reported that all patients, except one, received transplants mismatched for one, two, or three antigens and that graft-versus-host disease (GVHD) was mild (or absent) in all but two recipients, who had grade III GVHD.

Recently, one of our patients, a five-year-old white boy with acute lymphoblastic leukemia in second complete remission, died of grade IV GVHD after receiving placental blood from an Asian donor, provided by the Dusseldorf Placental Blood Bank in Germany. Patient and donor were identical for the class I HLA-A, B, and C loci and class II HLA-DR, with a minor mismatch at the HLA-DQB1 locus (HLA-DQB1\*0602 in the case of the patient and HLA-DQB1\*0603 in the case of placental blood). Cyclosporine alone was used for GVHD prophylaxis. Although Kurtzberg et al. specify the racial and ethnic origins of their patients, they do not do so for the donors. It would be interesting to know whether Patients 13 and 16, who were mismatched for one and two antigens, respectively, were given placental blood from racially or ethnically different donors. If that was the case, race or ethnic group may have to be taken into account when placental-blood transplantation is being considered.

Manuel M. Abecasis, M.D.

António Guimaraes, M.D.

Alexandra Machado, M.D.

*Instituto Portugues de Oncologia*

*1930 Lisbon, Portugal*

The authors reply:

*To the Editor:* We agree that manipulation of the infant during delivery could affect the amount of placental blood collected for banking and transplantation. However, we stress that the placental-blood units banked at the New York Blood Center are collected from the delivered placenta, ex vivo, by a dedicated team whose members have no role in the care of the infant or mother. All units are screened for blood-borne pathogens, including hepatitis B and C virus, human T-cell lymphotropic virus types I and II, HIV, and cytomegalovirus, as well as for hemoglobinopathies. Test results are forwarded to the mother's and infant's physicians.

We agree that children should benefit from this research, and indeed, they do. The majority of transplantations have been performed in children: 44 percent of the 272 patients who have received placental-blood units from the New York Blood Center have been under 6 years of age, and another 29 percent were between the ages of 6 and 11 years. We believe that the blood of the infant donor, which previously was discarded, should be a public resource and that its use should involve no financial gain to any party.

Drs. Bertolini and Robustelli della Cuna question the CD34 cell counts we reported. The numbers are correct, but they represent the total CD34 population, not an immature (e.g., CD34+CD38-) subgroup. We have not observed a correlation between the number of CD34 cells infused and the recovery of the white-cell, platelet, or red-cell count.

To respond to Locatelli et al., we reviewed our information regarding reconstitution of B cells in our patients. Our results agree, in part, with their observation; approximately 50 to 75 percent of our patients had elevated percentages of CD19+ and CD20+ cells 6 to 12 months after transplantation, but no increases in the absolute numbers of B lymphocytes in 27 patients tested at 3 to 6 months and 10 patients tested at >12 months. Since all our patients were given prophylactic intravenous immune globulin during the first year after transplantation, we cannot comment on immunoglobulin production.

To Abecasis et al. we reply that Patient 13 was of Native American and black heritage, and her donor was of white and black heritage. Both Patient 16 and the donor were white.

Finally, we would like to point out that the key to Figure 1 is reversed: 22 patients received placental blood thawed with the new method, and 3 patients received placental blood thawed with the old method.

Joanne Kurtzberg, M.D.

*Duke University Medical Center*

*Durham, NC 27710*

Pablo Rubenstein, M.D.

Cladd E. Stevens, M.D.

*New York Blood Center*

*New York, NY 10021*

*To the Editor:* Dr. Ammann is concerned that adults will benefit more than infants and children from placental-blood banking, that parents of donors may not be informed of positive results of tests for genetic

or infectious diseases, that the rights of infants as donors will be ignored, and that placental-blood donors may be rendered anemic by the process.

We agree that institutions that propose to collect and store placental blood need to address important medical and ethical issues. It is important to establish guidelines and regulations to ensure medical safety and protect the rights of placental-blood donors and potential recipients. Adherence to such regulations has been standard in the process of volunteer blood donation in this country for decades. We endorse the suggestion that the Food and Drug Administration or another appropriate organization should be responsible for establishing and enforcing similar standards for placental-blood collection.

In all but a very few cases thus far, children rather than adults have received placental blood. It has not been shown that placental blood consistently engrafts in adult-sized recipients, but recipients of placental blood should be selected on the basis of medical need and HLA compatibility. Placental-blood banking will thus benefit adults and children alike. We do not condone the development of commercial banks for allogeneic or autologous banking. On the contrary, we support the concept that in cases in which placental blood would otherwise be discarded (e.g., the great majority of deliveries), this blood represents a commonly owned good that, if collected for potential future transplantation, should be maintained and distributed by publicly accessible banks.

Other ethical issues regarding placental-blood banking include some of those raised by Dr. Ammann. These relate to informed consent for the collection and storage of placental blood, the obligation to notify parents and children of the results of tests performed on placental blood, the assurance of the confidentiality of such information, and the need for equitable access to placental blood.<sup>1</sup> Published policies and procedures proposed for placental-blood banks address these issues and would require, for example, that the donor be notified of test results.<sup>2</sup>

The timing of umbilical-cord clamping at delivery and the position of the infant relative to the placenta affect the volume of blood transfused to the infant and the volume remaining in the umbilical cord and placenta. The availability of placental blood for autotransfusion is of concern primarily in premature infants, who might require emergency resuscitation at delivery or who may benefit from later administration of additional blood.<sup>3,4</sup> Proposed policies for placental-blood banks restrict collection to full-term, uncomplicated pregnancies.<sup>2</sup> Moreover, in our institution and others, it is stipulated that placental-blood collection cannot interfere with the usual obstetrical practice at delivery and that cord clamping proceed accordingly.

Leigh C. Jefferies, M.D.

Leslie E. Silberstein, M.D.

*University of Pennsylvania School of Medicine  
Philadelphia, PA 19104*

## References

1. Sugarman J, Reisner EG, Kurtzberg J. Ethical aspects of banking placental blood for transplantation. *JAMA* 1995;274:1783-1785.
2. McCullough J, Clay ME, Fautsch S, et al. Proposed policies and procedures for the establishment of a cord blood bank. *Blood Cells* 1994;20:609-626.
3. Ballin A, Arbel E, Kenet G, et al. Autologous umbilical cord blood transfusion. *Arch Dis Child* 1995;73:F181-F183.
4. Kinmond S, Aitchison TC, Holland BM, Jones JG, Turner TL, Wardrop CA. Umbilical cord clamping and preterm infants: a randomised trial. *BMJ* 1993;306:172-175.