

# The New England Journal of Medicine

Copyright © 2002 by the Massachusetts Medical Society

VOLUME 346

JANUARY 17, 2002

NUMBER 3



## OUTCOMES IN YOUNG ADULTHOOD FOR VERY-LOW-BIRTH-WEIGHT INFANTS

MAUREEN HACK, M.B., CH.B., DANIEL J. FLANNERY, PH.D., MARK SCHLUCHTER, PH.D., LYDIA CARTAR, M.A.,  
ELAINE BORAWSKI, PH.D., AND NANCY KLEIN, PH.D.

### ABSTRACT

**Background** Very-low-birth-weight infants (those weighing less than 1500 g) born during the initial years of neonatal intensive care have now reached young adulthood.

**Methods** We compared a cohort of 242 survivors among very-low-birth-weight infants born between 1977 and 1979 (mean birth weight, 1179 g; mean gestational age at birth, 29.7 weeks) with 233 controls from the same population in Cleveland who had normal birth weights. We assessed the level of education, cognitive and academic achievement, and rates of chronic illness and risk-taking behavior at 20 years of age. Outcomes were adjusted for sex and socio-demographic status.

**Results** Fewer very-low-birth-weight young adults than normal-birth-weight young adults had graduated from high school (74 percent vs. 83 percent,  $P=0.04$ ). Very-low-birth-weight men, but not women, were significantly less likely than normal-birth-weight controls to be enrolled in postsecondary study (30 percent vs. 53 percent,  $P=0.002$ ). Very-low-birth-weight participants had a lower mean IQ (87 vs. 92) and lower academic achievement scores ( $P<0.001$  for both comparisons). They had higher rates of neurosensory impairments (10 percent vs. <1 percent,  $P<0.001$ ) and subnormal height (10 percent vs. 5 percent,  $P=0.04$ ). The very-low-birth-weight group reported less alcohol and drug use and had lower rates of pregnancy than normal-birth-weight controls; these differences persisted when comparisons were restricted to the participants without neurosensory impairment.

**Conclusions** Educational disadvantage associated with very low birth weight persists into early adulthood. (N Engl J Med 2002;346:149-57.)

Copyright © 2002 Massachusetts Medical Society.

THE introduction of neonatal intensive care in the 1960s resulted in substantial improvements in outcomes for very-low-birth-weight infants (those weighing less than 1500 g). By the 1970s, 80 to 90 percent of survivors were reported to be free of serious handicaps.<sup>1</sup> However, at school age, very-low-birth-weight children have poorer cognitive function and academic performance than normal-birth-weight controls.<sup>2-5</sup> Learning problems at school persist into adolescence and are apparent even in children who have normal intelligence and no neurologic impairment.<sup>6,7</sup>

Before the era of neonatal intensive care, when few very-low-birth-weight infants survived, the majority of survivors were described as having average or above-average mental development.<sup>8</sup> A more recent report, however, indicated that fewer persons with birth weights of less than 1000 g than normal-birth-weight controls had graduated from high school.<sup>9</sup> Reports from Europe note that very-low-birth-weight young adults have similar levels of educational attainment<sup>10,11</sup> but more chronic illness and handicap<sup>11-13</sup> than members of the general population<sup>10,12,13</sup> or normal-birth-weight controls.<sup>11</sup>

We undertook a longitudinal study of very-low-birth-weight children born in 1977, 1978, or 1979, and we previously reported on outcomes at eight years of age.<sup>2,14-17</sup> The current report extends the follow-up to 20 years of age. We hypothesized that, as compared with normal-birth-weight controls, very-low-birth-weight young adults would have lower intelligence, lower levels of educational achievement, and

From the Departments of Pediatrics (M.H., D.J.F., M.S., L.C.) and Epidemiology and Biostatistics (E.B.), Case Western Reserve University, and the Department of Special Education, Cleveland State University (N.K.) — both in Cleveland; and the Institute for the Study and Prevention of Violence, Kent State University, Kent, Ohio (D.J.F.). Address reprint requests to Dr. Hack at the Rainbow Babies and Children's Hospital of the University Hospitals of Cleveland, 11100 Euclid Ave., Cleveland, OH 44106, or at mxh7@po.cwru.edu.

higher rates of chronic health conditions and risk-taking behavior.

## METHODS

### Very-Low-Birth-Weight Group

A cohort of 490 very-low-birth-weight infants were admitted to Rainbow Babies and Children's Hospital in Cleveland between 1977 and 1979. A total of 316 children (64 percent) survived to their second year. One child died of a brain tumor between 2 and 8 years of age, and three died between 8 and 20 years of age — one from meningitis, one by drowning, and one from sequelae of severe spastic quadriplegia. Of the remaining 312 subjects, 70 were not studied: 58 could not be located, 5 lived out of state, 6 declined to participate, and 1 with severe spastic quadriplegia could not be interviewed. The study population thus included 242 very-low-birth-weight participants; this population represented 78 percent of the surviving birth cohort and included 232 (91 percent) of the 256 members of the cohort who had participated at eight years of age and an additional 10 who had not participated at eight years of age. One parent of each of 227 of the study participants (94 percent) was also interviewed. The mother was the parent interviewed in 91 percent of the cases.

The 242 very-low-birth-weight participants had a mean birth weight of 1179 g and had been born at a mean gestational age of 29.7 weeks; 110 (45 percent) had been born at the adjacent perinatal center of MacDonal Hospital for Women. Fifty-eight (24 percent) had a birth weight of less than 1000 g. Neonatal problems had included respiratory distress syndrome in 181 infants (75 percent), apnea of prematurity in 123 (51 percent), sepsis in 34 (14 percent), and necrotizing enterocolitis in 9 (4 percent). Assisted ventilation had been provided to 107 of the infants (44 percent). There were no major congenital malformations or congenital infections. The cohort was born before the advent of cerebral ultrasonography; thus, the rates of periventricular hemorrhage and leukomalacia are unknown. At the time of their birth, the mean age of their mothers was 24 years. A total of 158 of the mothers (65 percent) were married, and 59 (24 percent) had less than a high-school education.

The very-low-birth-weight young adults who participated in the study did not differ significantly from those who had been lost to follow-up since birth in terms of the sociodemographic characteristics of their mothers at the time of their birth: 35 percent of the mothers of participants were unmarried, as compared with 39 percent of the mothers of those lost to follow-up; 55 percent of the mothers were black, as compared with 46 percent among those lost to follow-up; and 24 percent had less than a high-school education, as compared with 33 percent among those lost to follow-up. More of those who participated at 20 years of age than of those who were lost to follow-up had been born at the perinatal center (46 percent vs. 31 percent,  $P=0.03$ ). However, the mean birth weight (1179 g vs. 1187 g), the mean gestational age at birth (29.7 weeks vs. 29.8 weeks), and the incidence of neonatal problems (including respiratory distress syndrome requiring ventilator therapy, sepsis, and necrotizing enterocolitis) did not differ between the groups. The mean IQ at 8 years of age was  $95 \pm 18$  among those who participated at 20 years of age and  $91 \pm 21$  among those who were subsequently lost to follow-up ( $P=0.27$ ).

### Control Group

The original control group included 366 normal-birth-weight children born at term in 1977, 1978, or 1979, who were selected by means of a population-sampling procedure when they were eight years of age.<sup>2</sup> Three of the controls died between 8 and 20 years of age — one from a gunshot wound, one from a stab wound, and one by suicide. Of the remaining 363 controls, 130 were not studied at 20 years of age: 91 could not be located, 1 lived out of state, and 38 declined to participate. The control population thus in-

cluded 233 participants — 64 percent of the cohort that had been recruited at eight years of age. One parent of each of 218 of the controls (94 percent) was also interviewed; in 95 percent of cases, the parent interviewed was the mother.

The normal-birth-weight controls who participated at 20 years of age had had a significantly higher mean IQ score on the Wechsler Intelligence Scale for Children-Revised (WISC-R) at 8 years of age than those who did not participate ( $104 \pm 17$  vs.  $93 \pm 14$ ). Fewer mothers of the controls who participated at 20 years of age than mothers of those who did not participate were unmarried when the child was 8 years old (36 percent vs. 61 percent), fewer had less than a high-school education (11 percent vs. 27 percent), and fewer were black (55 percent vs. 76 percent;  $P<0.001$  for all comparisons).

### Measures at 20 Years of Age

All subjects provided written informed consent to participate in the study. Information concerning educational attainment, current enrollment in an educational program, and other activities was obtained from the young adults by means of interviews. High-school graduation was confirmed by means of school records. Intelligence was assessed with the Short Form of the Wechsler Adult Intelligence Scale-Revised (WAIS-R); we used the Vocabulary and Block Design subtests, which measure verbal comprehension and perceptual-organization skills, respectively.<sup>18,19</sup> Academic skills were assessed with the Letter-Word Identification and Applied Problems subtests of the Tests of Achievement from the Woodcock-Johnson Psycho-Educational Battery-Revised.<sup>20</sup> Health status was ascertained from the participants by means of questions concerning chronic medical, neurologic, or psychiatric conditions that had lasted 12 months or longer, as well as detailed questions concerning pregnancy and childbirth.<sup>21,22</sup> Height was measured with a Harpenden stadiometer. The extent of engagement in risk-taking behavior during the previous 12 months was ascertained by means of self-administered questionnaires — a substance-abuse checklist and a Sexual Experience Scale<sup>23,24</sup> — with additional questions concerning contact with the police. The subjects were asked whether they had ever been in jail, including for several hours or overnight or in juvenile detention, and whether they had been convicted of a crime, including driving under the influence of alcohol. The parents completed questionnaires regarding the young adults' health<sup>21</sup> and their knowledge about substance use by the participants.<sup>23</sup>

### Statistical Analysis

Univariate comparisons between the very-low-birth-weight and normal-birth-weight groups were made with the use of Student's *t*-test for continuous variables and with the chi-square test or Fisher's exact test for discrete variables. Logistic regression was used for dichotomous outcomes, and multiple linear regression was used for continuous outcomes. Because of the effects of sociodemographic factors on outcomes, we controlled for sociodemographic status in the analyses.<sup>2,4,6</sup> Because of differences in behavior between the sexes, we performed separate analyses for each sex, comparing the groups after adjusting for sociodemographic status. We also compared the groups using pooled data from participants of both sexes with adjustment for sex and sociodemographic status. If the interaction between birth-weight groups and sex was found to be significant, indicating that the effect on the outcome differed between male participants and female participants, then the adjusted difference from the pooled analysis was not presented. A composite score representing the mother's sociodemographic status, which we used previously,<sup>2</sup> was calculated by assigning one point for each of the following factors: unmarried status, black race, and less than a high-school education.<sup>2</sup> The composite score ranged from zero to three. We used the mother's marital and educational status at the time the child was eight years old, since it was considered more relevant than her later status to the period of child de-

velopment that is critical for educational attainment. This composite had a stronger correlation with the IQ at 8 years of age<sup>2</sup> and at 20 years of age than did any of its components analyzed separately. In separate sets of analyses, we compared the subgroups of very-low-birth-weight and normal-birth-weight participants who did not have neurosensory impairment, subnormal IQ, or chronic conditions.

RESULTS

Sociodemographic Status and Chronic Health Conditions

The very-low-birth-weight and normal-birth-weight groups did not differ significantly with regard to maternal marital status at eight years of follow-up or maternal race. The level of maternal education was lower in the very-low-birth-weight group, but the composite index of sociodemographic status did not differ significantly between groups (Table 1). Six of the very-low-birth-weight young adults and one of the normal-birth-weight young adults were married.

Very-low-birth-weight participants had significantly higher rates of chronic conditions than the controls (33 percent vs. 21 percent,  $P=0.002$ ). The difference was primarily attributable to higher rates of neurosensory impairment and subnormal height<sup>25</sup> (Table 2). A total of 23 percent of the very-low-birth-weight participants had one chronic condition, 9 percent had two chronic conditions, and 1 percent had three or more chronic conditions. In comparison, 17 percent of the controls had one chronic condition, and 4 percent had two chronic conditions ( $P=0.005$ ).

Educational Attainment and Current Enrollment in an Educational Program

Fewer very-low-birth-weight participants than normal-birth-weight participants had graduated from high school or obtained a general equivalency diploma by 20 years of age (74 percent vs. 83 percent,  $P=0.04$ ) (Table 3). Forty percent of the very-low-birth-weight participants had repeated a grade in school, as compared with 27 percent of the normal-birth-weight participants ( $P=0.003$ ). Very-low-birth-weight participants who graduated from high school did so at a mean age of  $18.2 \pm 0.6$  years, as compared with  $17.9 \pm 0.6$  years among the controls ( $P<0.001$ ). Fewer very-low-birth-weight men were enrolled in postsecondary studies, of whom only 16 percent were in a four-year college program, as compared with 44 percent in the control group ( $P<0.001$ ).

The differences in grade repetition, educational attainment, and current enrollment in educational programs remained significant when the comparisons were restricted to participants without neurosensory impairment or subnormal IQ ( $<70$ ).

Intelligence and Academic Achievement

Very-low-birth-weight participants had significantly lower mean IQ scores than the controls (87 vs. 92,

TABLE 1. MATERNAL SOCIODEMOGRAPHIC STATUS AT EIGHT YEARS OF FOLLOW-UP AND BIRTH DATA FOR VERY-LOW-BIRTH-WEIGHT AND NORMAL-BIRTH-WEIGHT PARTICIPANTS.\*

VARIABLE	VERY LOW BIRTH WEIGHT (N=242)	NORMAL BIRTH WEIGHT (N=233)
<b>Maternal characteristics — no. (%)</b>		
Unmarried	100 (41)	84 (36)
Black race	133 (55)	128 (55)
Education†		
<High-school graduation	40 (17)	25 (11)
High-school graduation	133 (55)	118 (51)
>High-school graduation	69 (29)	90 (39)
Composite sociodemographic score‡		
0	79 (33)	90 (39)
1	73 (30)	66 (28)
2	71 (29)	60 (26)
3	19 (8)	17 (7)
<b>Birth data</b>		
Weight — g	1179 ± 219	3279 ± 584
Gestational age — wk	29.7 ± 2	≥ 37§
Female sex — no. (%)	126 (52)	125 (54)
Multiple birth — no. (%)¶	32 (13)	4 (2)

\*Plus-minus values are means ± SD. The maternal status at the time the participant was eight years old is given. Maternal characteristics refer to participant's primary caretaker, who was an adoptive mother in five instances in the very-low-birth-weight group and three instances in the control group, a foster mother in one instance in the very-low-birth-weight group, and a grandparent in three instances in the very-low-birth-weight group.

† $P=0.03$  for the comparison between groups.

‡In the calculation of this composite score, one point was assigned for each of the following factors: unmarried status, black race, and less than a high-school education.

§Specific information on gestational age was not available for the control group.

¶Data are for participants with a living twin or, in one case, two living triplets.

$P<0.001$ ) and had lower scores on the subtests of academic achievement (Table 4).<sup>20</sup> They also had a higher frequency of subnormal IQ ( $<70$ ) and borderline IQ (70 to 84). Fifty-one percent of the very-low-birth-weight participants had an IQ in the normal range ( $\geq 85$ ), as compared with 67 percent of the controls ( $P<0.001$ ). These differences remained significant when the comparisons were restricted to the participants without neurosensory impairment. There was a significant interaction of birth weight and sex in the scores on the Applied Problems subtest of the Woodcock-Johnson Psycho-Educational Battery-Revised, with a greater difference between groups for male participants than for female participants.

Substance Use, Contact with Police, and Sexual Activity

The rates of smoking did not differ significantly between the groups, but the very-low-birth-weight participants reported significantly lower rates of alcohol

**TABLE 2. CHRONIC CONDITIONS AT 20 YEARS OF AGE AMONG VERY-LOW-BIRTH-WEIGHT AND NORMAL-BIRTH-WEIGHT PARTICIPANTS.\***

VARIABLE	MEN		WOMEN	
	VERY LOW BIRTH WEIGHT (N=116)	NORMAL BIRTH WEIGHT (N=108)	VERY LOW BIRTH WEIGHT (N=126)	NORMAL BIRTH WEIGHT (N=125)
	no. of participants (%)			
Neurosensory condition	11 (9)	1 (1)†	14 (11)	0‡
Cerebral palsy§	6 (5)	0	9 (7)	0
Hydrocephalus necessitating the placement of a shunt	1 (1)	0	4 (3)	0
Blindness¶	3 (3)	0	1 (1)	0
Deafness	1 (1)	1 (1)	2 (2)	0
Medical or psychiatric illness	22 (19)	17 (16)	29 (23)	20 (16)
Asthma**	8 (7)	6 (6)	11 (9)	7 (6)
Diabetes	0	0	1 (1)	1 (1)
Sickle cell anemia	0	1 (1)	1 (1)	1 (1)
Epilepsy	1 (1)	1 (1)	3 (2)	0
Arthritis	3 (3)	2 (2)	10 (8)	6 (5)
Bone or muscle disorder	10 (9)	7 (6)	5 (4)	7 (6)
Bipolar disorder	3 (3)	2 (2)	2 (2)	0
Other††	0	0	1 (1)	2 (2)
Height <3rd percentile‡‡	9 (8)	5 (5)	14 (11)	6 (5)
Total with at least one condition	36 (31)	23 (21)	45 (36)	25 (20)§§

\*Chronic conditions were defined as those with a duration of 12 months or more. Data for the general categories of neurosensory condition and medical or psychiatric illness are the numbers and percentages of participants with at least one condition in that category.

†P=0.004 for the comparison with the men in the very-low-birth-weight group.

‡P=0.005 for the comparison with the women in the very-low-birth-weight group.

§Nine of the participants had spastic diplegia, two had spastic hemiplegia, and four had spastic quadriplegia.

¶One participant had bilateral blindness, and three had unilateral blindness.

||Data are for participants who required a hearing aid.

\*\*Data are for participants who had had an asthma attack in the previous 12 months, were taking asthma medication, or both.

††One participant in the very-low-birth-weight group had hypertension due to Liddle's syndrome (pseudoaldosteronism); one participant in the control group had endometriosis, and one had narcolepsy.

‡‡Height percentiles from the Centers for Disease Control and Prevention growth charts<sup>25</sup> were used. The analysis includes 11 participants who were not measured but reported their own height.

§§P=0.006 for the comparison with the women in the very-low-birth-weight group.

and marijuana use than the controls — differences primarily attributable to differences among the women (Table 5). These differences remained significant when the comparisons were restricted to participants without neurosensory impairment, subnormal IQ, or chronic conditions. Parents also reported significantly lower rates of alcohol use for very-low-birth-weight participants but similar rates of smoking, marijuana use, and other illicit drug use (data not shown).

Fewer very-low-birth-weight men than male controls had ever had contact with the police — a difference that was primarily attributable to less contact related to truancy (5 percent vs. 14 percent, P=0.03) and less contact related to drug or alcohol use (13 percent vs. 29 percent, P=0.006). Fewer very-low-birth-

weight women than female controls reported ever having intercourse, being pregnant, or delivering a live-born infant (Table 5). The differences between the groups in the rates of substance abuse, contact with the police, and pregnancy remained significant when the comparisons were restricted to participants without neurosensory impairment or subnormal IQ. There were no significant differences between the very-low-birth-weight group and the normal-birth-weight group in the rates of conviction for a crime or incarceration in jail or juvenile detention.

## DISCUSSION

Fewer very-low-birth-weight participants than normal-birth-weight participants had graduated from high

**TABLE 3. LEVEL OF EDUCATION AT 20 YEARS OF AGE AMONG VERY-LOW-BIRTH-WEIGHT AND NORMAL-BIRTH-WEIGHT PARTICIPANTS.\***

VARIABLE	MEN			WOMEN			TOTAL POPULATION
	VERY LOW BIRTH WEIGHT (N=116)	NORMAL BIRTH WEIGHT (N=108)	ODDS RATIO (95% CI)	VERY LOW BIRTH WEIGHT (N=126)	NORMAL BIRTH WEIGHT (N=125)	ODDS RATIO (95% CI)	ODDS RATIO (95% CI)
	no. (%)			no. (%)			
High-school graduation†	77 (66)	81 (75)	0.7 (0.4–1.3)	102 (81)	112 (90)	0.5 (0.2–1.1)	0.6 (0.4–1.0)‡
Current study							
None	70 (60)	44 (41)	2.1 (1.2–3.7)§	56 (44)	53 (42)	1.1 (0.6–1.8)	1.5 (1.0–2.1)‡
High school or GED¶	11 (9)	8 (7)	1.2 (0.5–3.2)	6 (5)	4 (3)	1.4 (0.4–5.3)	1.3 (0.6–2.8)
Postsecondary study	35 (30)	57 (53)	0.4 (0.2–0.7)**	64 (51)	68 (54)	0.9 (0.5–1.5)	
Community college††	17 (15)	9 (8)	1.9 (0.8–4.5)	22 (17)	21 (17)	1.0 (0.5–2.0)	1.3 (0.8–2.2)
Four-year college‡‡	18 (16)	47 (44)	0.2 (0.1–0.4)§§	42 (33)	47 (38)	0.8 (0.5–1.5)	

\*The odds ratios for men and for women were adjusted for sociodemographic status; the odds ratios for the total population were adjusted for sociodemographic status and sex. CI denotes confidence interval, and GED general equivalency diploma.

†Data include 12 participants in the very-low-birth-weight group and 17 controls who had obtained a GED.

‡P=0.04 for the comparison between groups.

§P=0.007 for the comparison between groups.

¶Twelve very-low-birth-weight participants and seven controls were in high school, and five very-low-birth-weight participants and five controls were in a GED program.

||P=0.04 for the interaction between birth weight and sex. Because this interaction was significant, the adjusted difference from the pooled analysis is not presented.

\*\*P=0.002 for the comparison between groups.

††Data include 2 very-low-birth-weight participants and 4 controls who were in business school and 15 very-low-birth-weight participants and 7 controls who were in technical school.

‡‡P=0.004 for the interaction between birth weight and sex. Because this interaction was significant, the adjusted difference from the pooled analysis is not presented.

§§P<0.001 for the comparison between groups.

school or obtained an equivalency diploma by 20 years of age. Very-low-birth-weight participants were less likely to be enrolled in a postsecondary educational program — a difference primarily attributable to the difference between the men in the two groups. The very-low-birth-weight participants had a higher incidence of chronic medical conditions, mainly because of higher rates of neurosensory impairment and subnormal height. Contrary to our expectations, the very-low-birth-weight group reported less risk-taking behavior than the control group, including less use of alcohol and illicit substances. Fewer very-low-birth-weight men than control men had ever had contact with the police, and fewer very-low-birth-weight women than control women had ever had intercourse, been pregnant, or had a baby.

Data have been lacking on the follow-up into adulthood of very-low-birth-weight survivors of neonatal intensive care in the United States. The very-low-birth-weight participants in our study had been treated in an urban perinatal center, and our study population

thus included more persons of lower socioeconomic status and minority race or ethnic group than the United States as a whole. The rates of chronic illness in our normal-birth-weight control population are similar to those reported for young adults nationally,<sup>21</sup> and their rates of substance abuse, contact with the police, and childbirth are in keeping with data from similar urban populations.<sup>26–30</sup>

We followed 78 percent of the birth cohort of very-low-birth-weight subjects to 20 years of age, but only 64 percent of the normal-birth-weight subjects enrolled at 8 years of age remained in the study until they were 20 years old. The loss of very-low-birth-weight subjects occurred mainly between birth and eight years of age. Those who were followed to 20 years of age were similar to the original birth cohort in terms of birth weight, gestational age at birth, and neonatal morbidity, although their mothers tended to have a higher level of education. The very-low-birth-weight and normal-birth-weight groups were similar in terms of maternal sociodemographic charac-

**TABLE 4. COGNITIVE TEST RESULTS AT 20 YEARS OF AGE AMONG VERY-LOW-BIRTH-WEIGHT AND NORMAL-BIRTH-WEIGHT PARTICIPANTS.\***

VARIABLE	MEN			WOMEN			TOTAL POPULATION			
	VERY LOW BIRTHWEIGHT (N=113)	NORMAL BIRTHWEIGHT (N=106)	P VALUE	VERY LOW BIRTHWEIGHT (N=123)	NORMAL BIRTHWEIGHT (N=125)	P VALUE	ODDS RATIO (95% CI)	P VALUE	ODDS RATIO (95% CI)	P VALUE
IQ range										
<70	9 (8)	2 (2)	4.6 (0.9 to 22.6)	7 (6)	2 (2)	3.6 (0.7 to 18.0)		4.0 (1.3 to 12.2)	0.02	
70-84	44 (39)	23 (22)	2.3 (1.2 to 4.3)	56 (46)	49 (39)	1.3 (0.8 to 2.2)		1.7 (1.1 to 2.5)	0.02	
≥85	60 (53)	81 (76)	0.3 (0.2 to 0.6)	60 (49)	74 (59)	0.6 (0.4 to 1.1)		0.5 (0.3 to 0.7)	<0.001	
	no. (%)	no. (%)		no. (%)	no. (%)					
IQ score										
Academic achievement scores										
Letter-Word Identification	87.6±15.1	94.7±14.9	-5.9 (-9.2 to -2.6)	86.2±13.4	89.8±14.0	-3.4 (-6.5 to -0.4)		-4.6 (-6.9 to -2.4)	<0.001	
Applied Problems†	94.4±19.8	103.2±20.2	-7.6 (-12.4 to -2.9)	97.1±19.2	102.3±21.7	-5.0 (-9.7 to -0.2)		-6.3 (-9.6 to -2.9)	<0.001	
	89.0±14.2	98.4±16.8	-8.0 (-11.6 to -4.5)	89.0±12.3	92.4±12.4	-3.1 (-6.0 to -0.3)				
	VERY LOW BIRTHWEIGHT	NORMAL BIRTHWEIGHT	DIFFERENCE (95% CI)	VERY LOW BIRTHWEIGHT	NORMAL BIRTHWEIGHT	DIFFERENCE (95% CI)				

\*The IQ was measured by the Short Form of the Wechsler Adult Intelligence Scale-Revised,<sup>18</sup> which includes the Vocabulary and Block Design subscales.<sup>18,19</sup> The standard score has a mean of 100 and a standard deviation of 15. Academic achievement was assessed by the Tests of Achievement from the Woodcock-Johnson Psycho-Educational Battery-Revised.<sup>20</sup> The standard score has a mean of 100 and a standard deviation of 15. Scores on the Letter-Word Identification subtest were available for 235 very-low-birth-weight participants and 228 controls; scores on the Applied Problems subtest were available for 238 very-low-birth-weight participants and 230 controls. Plus-minus values are means ±SD. The values shown are unadjusted. The odds ratios for men and for women were adjusted for sociodemographic status; the odds ratios for the total population were adjusted for sociodemographic status and sex. The differences in score between groups were calculated by first adjusting the mean values among men and among women for sociodemographic status and adjusting the mean values in the total population for sociodemographic status and sex and then subtracting the adjusted mean value for normal-birth-weight participants from the adjusted mean value for very-low-birth-weight participants. CI denotes confidence interval.

†P=0.02 for the interaction between birth weight and sex. Because this interaction was significant, the adjusted difference from the pooled analysis is not presented.

OUTCOMES IN YOUNG ADULTHOOD FOR VERY-LOW-BIRTH-WEIGHT INFANTS

TABLE 5. SELF-REPORTED SUBSTANCE USE, CRIMINAL ACTIVITY, AND SEXUAL ACTIVITY AT 20 YEARS OF AGE AMONG VERY-LOW-BIRTH-WEIGHT AND NORMAL-BIRTH-WEIGHT PARTICIPANTS.\*

VARIABLE	MEN				WOMEN				TOTAL POPULATION	
	VERY LOW BIRTH WEIGHT (N=116)	NORMAL BIRTH (N=108)	ODDS RATIO (95% CI)	P VALUE	VERY LOW BIRTH WEIGHT (N=126)	NORMAL BIRTH (N=124)	ODDS RATIO (95% CI)	P VALUE	ODDS RATIO (95% CI)	P VALUE
	no. (%)				no. (%)					
Substance use during the previous year										
Tobacco	66 (57)	64 (59)	0.9 (0.5–1.6)		50 (40)	59 (48)	0.7 (0.4–1.2)		0.8 (0.6–1.2)	
Alcohol	84 (72)	89 (82)	0.6 (0.3–1.1)		77 (61)	103 (83)	0.3 (0.2–0.6)	<0.001	0.4 (0.3–0.6)	<0.001
Illicit drugs	49 (42)	57 (53)	0.6 (0.4–1.1)		38 (30)	54 (44)	0.6 (0.3–0.9)	0.03	0.6 (0.4–0.9)	0.007
Marijuana	49 (42)	56 (52)	0.7 (0.4–1.1)		37 (29)	52 (42)	0.6 (0.3–1.0)	0.04	0.6 (0.4–0.9)	0.01
Other†	10 (9)	9 (8)	1.1 (0.4–2.9)		5 (4)	9 (7)	0.5 (0.2–1.7)		0.8 (0.4–1.7)	
Contact with the police										
Violation of law (excluding traffic laws)	43 (37)	56 (52)	0.5 (0.3–0.9)	0.03	30 (24)	29 (23)	1.0 (0.6–1.8)		0.7 (0.5–1.1)	
Convicted of crime‡	23 (20)	29 (27)	0.7 (0.4–1.2)		3 (2)	4 (3)	0.7 (0.1–3.2)		0.7 (0.4–1.2)	
Incarcerated§	30 (26)	28 (26)	0.9 (0.5–1.7)		8 (6)	7 (6)	1.1 (0.4–3.2)		0.9 (0.6–1.6)	
Sexual activity¶										
Intercourse	96 (83)	88 (81)	0.8 (0.4–1.7)		82 (65)	97 (78)	0.5 (0.2–0.8)	0.01	0.6 (0.4–0.9)	0.02
Pregnancy	30 (26)	25 (23)	1.1 (0.6–2.0)		36 (29)	51 (41)	0.5 (0.3–0.9)	0.02	0.7 (0.5–1.1)	
Live birth**	17 (15)	19 (18)	0.7 (0.4–1.5)		17 (13)	30 (24)	0.4 (0.2–0.9)	0.02	0.6 (0.3–0.9)	0.02

\*The odds ratios for men and for women were adjusted for sociodemographic status; the odds ratios for the total population were adjusted for sociodemographic status and sex. CI denotes confidence interval.

†Data include the use of inhalants (by four very-low-birth-weight men, one male control, and one female control), amphetamines (by five very-low-birth-weight men, three very-low-birth weight women, three male controls, and five female controls), cocaine (by two very-low-birth-weight men, one very-low-birth-weight woman, two male controls, and three female controls), and hallucinogens (by three very-low-birth-weight men, three very-low-birth-weight women, seven male controls, and seven female controls).

‡Data include convictions for driving under the influence of alcohol.

§Incarceration was defined as ever being held in jail, including for several hours or overnight, or in juvenile detention.

¶For men, the data for pregnancy or live birth indicate pregnancy in the man's partner or live birth of a child fathered by the man.

||Data include 13 very-low-birth-weight women and 9 female controls who were pregnant at the time of the interview.

\*\*Data include three very-low-birth-weight women and five female controls who had had more than one live birth.

teristics at eight years of age, when the normal-birth-weight group was recruited.<sup>2</sup> From 8 to 20 years of age, both groups had greater losses to follow-up among children whose mothers had less education, but more of these losses occurred in the control group than in the very-low-birth-weight group. This imbalance explains the discrepancy in maternal education between the very-low-birth-weight participants and the controls at 20 years of age. Previous studies of very-low-birth-weight infants have similarly had greater loss to follow-up among participants with lower IQs and those whose mothers had lower levels of education and were of lower social class.<sup>31,32</sup> To control for this bias, we adjusted for maternal sociodemographic status, which included maternal educational level, in all the analyses. A weakness of the study is that we did not measure maternal IQ, an important predictor of children's educational outcomes.<sup>33</sup>

The findings that fewer very-low-birth-weight par-

ticipants than normal-birth-weight participants had graduated from high school and that they were less likely to be enrolled in a postsecondary educational program are not surprising, considering their lower IQs and academic achievement scores and higher rates of grade repetition. A disadvantage in school performance among boys has been previously reported in studies of very-low-birth-weight subjects.<sup>34,35</sup> Our finding of a lower rate of enrollment in four-year colleges among the very-low-birth-weight men, even after sociodemographic factors had been controlled for, indicates that this disadvantage extends into young adulthood. These findings suggest that men who had very low birth weight will lag behind their normal-birth-weight peers in their ultimate educational and occupational achievement, and thus in earning ability, social status, and prestige.<sup>36</sup>

Lower rates of alcohol and drug use among very-low-birth-weight subjects have not previously been re-

ported. Bjerager et al. noted similar rates of alcohol and drug use for very-low-birth-weight young adults and controls.<sup>11</sup> Chilcoat and Breslau reported an increase in early drug use among 11-year-old low-birth-weight children that was associated with attention-deficit-hyperactivity disorder rather than with birth weight per se.<sup>37</sup> Risk-taking behavior in children is associated with externalizing behavior,<sup>37</sup> low intelligence and academic performance,<sup>38</sup> negative peer influences,<sup>39</sup> early puberty,<sup>24</sup> poor self-esteem, and poor parental monitoring.<sup>23,40-42</sup> We do not have information on all of these correlated factors, but we postulate that the more limited risk-taking behavior that we have documented may result from increased parental monitoring of very-low-birth-weight children.

A problem inherent in the long-term follow-up of preterm infants is that outcomes might not be relevant to survivors of current neonatal intensive care. Recent survivors who weighed less than 1000 g at birth have poorer outcomes than were previously reported.<sup>43,44</sup> There is, however, no evidence that the incidence of neurodevelopmental sequelae of very low birth weight among children who weighed between 1000 and 1500 g at birth has changed since the late 1970s.<sup>45</sup> The majority of our 20-year-old cohort had birth weights in this range. We thus suggest that our results have relevance to current survivors with birth weights in the same range.

In summary, the results of this study indicate that the neurodevelopmental and growth-related sequelae of very low birth weight and the poor school achievement that have been reported for very-low-birth-weight children persist into young adulthood. The results are reassuring, however, in that these problems are not associated with increased risk-taking behavior or criminal activity. Further follow-up will be important to examine the ultimate educational attainment and choice of occupation of the cohort as they reach mature adulthood.

Supported by grants (RO1 HD34177 and General Clinical Research Center grant M01 RR00080) from the National Institutes of Health and by a grant (96-46) from the Genentech Foundation for Growth and Development. The latter did not participate in the study design, data collection, data analysis, or manuscript preparation.

*We are indebted to Debra Hoffman, Anne Rivers, Susan McGrath, Miriam Curran, Terry Reid, and Elizabeth Carter for their assistance in compiling and analyzing the data, and to Drs. A.A. Fanaroff, D. Drotar, and H.G. Taylor for their critical comments on the manuscript.*

## REFERENCES

- U.S. Congress, Office of Technology Assessment. Neonatal intensive care for low birthweight infants: costs and effectiveness (health technology case study 38). Washington, D.C.: Government Printing Office, December 1987. (OTA publication no. OTA-HCS-38.)
- Hack M, Breslau N, Aram D, Weissman B, Klein N, Borawski-Clark E. The effect of very low birth weight and social risk on neurocognitive abilities at school age. *J Dev Behav Pediatr* 1992;13:412-20.
- Botting N, Powlis A, Cooke RWI, Marlow N. Attention deficit hyperactivity disorders and other psychiatric outcomes in very low birthweight children at 12 years. *J Child Psychol Psychiatry* 1997;38:931-41.
- Saigal S. Follow-up of very low birthweight babies to adolescence. *Semin Neonatol* 2000;5:107-18.
- McCormick MC, Brooks-Gunn J, Workman-Daniels K, Turner J, Peckham GJ. The health and developmental status of very low-birth-weight children at school age. *JAMA* 1992;267:2204-8.
- Hack M, Klein NK, Taylor HG. Long-term developmental outcomes of low birth weight infants. *Future Child* 1995;5:176-96.
- Saigal S, Hoult LA, Streiner DL, Stoskopf B, Rosenbaum PL. School difficulties at adolescence in a regional cohort of children who were extremely low birth weight. *Pediatrics* 2000;105:325-31.
- Hess JH. Experiences gained in a thirty year study of prematurely born infants. *Pediatrics* 1953;11:425-34.
- Mazurier E, Lefebvre F, Tessier R. Educational achievement and intelligence at 16-21 years of ex-prematures born at  $\leq 1000$  g. *Pediatr Res* 1999;45:250A. abstract.
- Dessens AB, Smolders-de Haas H, Koppe JG. Twenty-year follow-up of antenatal corticosteroid treatment. *Pediatrics* 2000;105:1325. abstract. (Also available at <http://www.pediatrics.org/cgi/content/full/105/6/e77>.)
- Bjerager M, Steensberg J, Greisen G. Quality of life among young adults born with very low birthweights. *Acta Paediatr* 1995;84:1339-43.
- Ericson A, Kallen B. Very low birthweight boys at the age of 19. *Arch Dis Child Fetal Neonatal Ed* 1998;78:F171-F174.
- Xu B, Rantakallio P, Järvelin M-R. Mortality and hospitalizations of 24-year-old members of the low-birthweight cohort in northern Finland. *Epidemiology* 1998;9:662-5.
- Hack M, Breslau N, Weissman B, Aram D, Klein N, Borawski E. Effect of very low birth weight and subnormal head size on cognitive abilities at school age. *N Engl J Med* 1991;325:231-7.
- Hack M, Weissman B, Breslau N, Klein N, Borawski-Clark E, Fanaroff AA. Health of very low birth weight children during their first eight years. *J Pediatr* 1993;122:887-92.
- Hack M, Weissman B, Borawski-Clark E. Catch-up growth during childhood among very low-birth-weight children. *Arch Pediatr Adolesc Med* 1996;150:1122-9.
- Aram DM, Hack M, Hawkins S, Weissman BM, Borawski-Clark E. Very-low-birthweight children and speech and language development. *J Speech Hear Res* 1991;34:1169-79.
- Wechsler D. WAIS-R manual: Wechsler Adult Intelligence Scale-revised. San Antonio, Tex.: Psychological Corporation, 1981.
- Sattler JM, Ryan JJ. Wechsler Adult Intelligence Scale-revised (WAIS-R). In: Sattler JM, ed. *Assessment of children*. 3rd ed. Rev. San Diego, Calif.: Jerome M. Sattler, 1992.
- Woodcock RW, Mather N. WJ-R tests of achievement: standard and supplemental batteries: examiner's manual. Chicago: Riverside Publishing, 1989.
- Adams PF, Marano MA. Current estimates from the National Health Interview Survey, 1994. Vital and health statistics. Series 10. No. 193. Hyattsville, Md.: National Center for Health Statistics, 1995. (DHHS publication no. (PHS) 96-1521.)
- Starfield B, Riley AW, Breen BF, et al. The adolescent Child Health and Illness Profile: a population-based measure of health. *Med Care* 1995;33:553-66.
- Flannery DJ, Vazsonyi AT, Torquati J, Fridrich A. Ethnic and gender differences in risk for early adolescent substance use. *J Youth Adolesc* 1994;23:195-213.
- Flannery D, Rowe D, Gulley B. Impact of pubertal status, timing, and age on adolescent sexual experience and delinquency. *J Adolesc Res* 1993;8:21-40.
- Kuczumarski RJ, Ogden CL, Grummer-Strawn LM, et al. CDC growth charts: United States. Advance data from vital and health statistics. No. 314. Hyattsville, Md.: National Center for Health Statistics, 2000. (DHHS publication no. (PHS) 2000-1250 0-0431.)
- Federal Interagency Forum on Child and Family Statistics. *America's children: key national indicators of well-being*. Washington, D.C.: Government Printing Office, 2000.
- MacKay AP, Fingerhut LA, Duran CR. *Adolescent health chartbook: health*. United States, 2000. Hyattsville, Md.: National Center for Health Statistics, 2000. (DHHS publication no. 00-1232-1.)
- Kann L, Kinchen SA, Williams BI, et al. Youth risk behavior surveillance — United States, 1999. *CDC Surveill Summ* 2000;49(SS-5):1-32.
- Hardy JB, Shapiro S, Mellits ED, et al. Self-sufficiency at ages 27 to 33 years: factors present between birth and 18 years that predict educational attainment among children born to inner-city families. *Pediatrics* 1997;99:80-7.

30. Health-risk behaviors among persons aged 12–21 years — United States, 1992. *MMWR Morb Mortal Wkly Rep* 1994;43:231-5.
31. Aylward GP, Hatcher RP, Stripp B, Gustafson NF, Leavitt LA. Who goes and who stays: subject loss in a multicenter, longitudinal follow-up study. *J Dev Behav Pediatr* 1985;6:3-8.
32. Wolke D, Sohne B, Ohrt B, Riegel K. Follow-up of preterm children: important to document dropouts. *Lancet* 1995;345:447.
33. Bacharach VR, Baumeister AA. Effects of maternal intelligence, marital status, income, and home environment on cognitive development of low birthweight infants. *J Pediatr Psychol* 1998;23:197-205.
34. Verloove-Vanhorick SP, Veen S, Ens-Dokkum MH, Schreuder AM, Brand R, Ruys JH. Sex difference in disability and handicap at five years of age in children born at very short gestation. *Pediatrics* 1994;93:576-9.
35. Johnson EO, Breslau N. Increased risk of learning disabilities in low birth weight boys at age 11 years. *Biol Psychiatry* 2000;47:490-500.
36. Moss N. The body politic and the power of socioeconomic status. *Am J Public Health* 1997;87:1411-3.
37. Chilcoat HD, Breslau N. Pathways from ADHD to early drug use. *J Am Acad Child Adolesc Psychiatry* 2000;38:1347-54.
38. Halpern CT, Joyner K, Udry JR, Suchindran C. Smart teens don't have sex (or kiss much either). *J Adolesc Health* 2000;26:213-25.
39. Belcher HME, Shinitzky HE. Substance abuse in children: prediction, protection, and prevention. *Arch Pediatr Adolesc Med* 1998;152:952-60.
40. Chilcoat HD, Dishion TJ, Anthony JC. Parent monitoring and the incidence of drug sampling in urban elementary school children. *Am J Epidemiol* 1995;141:25-31.
41. Simantov E, Schoen C, Klein J. Health-compromising behaviors: why do adolescents smoke or drink? Identifying underlying risk and protective factors. *Arch Pediatr Adolesc Med* 2000;154:1025-33.
42. Flannery DJ, Williams LL, Vazsonyi AT. Who are they with and what are they doing? Delinquent behavior, substance use, and early adolescents' after-school time. *Am J Orthopsychiatry* 1999;69:247-53.
43. Hack M, Wilson-Costello D, Friedman H, Taylor GH, Schluchter M, Fanaroff AA. Neurodevelopment and predictors of outcomes of children with birth weights of less than 1000 g: 1992-1995. *Arch Pediatr Adolesc Med* 2000;154:725-31.
44. Vohr BR, Wright LL, Dusick AM, et al. Neurodevelopmental and functional outcomes of extremely low birth weight infants in the National Institute of Child Health and Human Development Neonatal Research Network, 1993-1994. *Pediatrics* 2000;105:1216-26.
45. Piccuch RE, Leonard CH, Cooper BA. Infants with birth weight 1,000-1,499 grams born in three time periods: has outcome changed over time? *Clin Pediatr (Phila)* 1998;37:537-45.

Copyright © 2002 Massachusetts Medical Society.

---

ELECTRONIC ACCESS TO THE *JOURNAL'S* CUMULATIVE INDEX

---

At the *Journal's* site on the World Wide Web (<http://www.nejm.org>) you can search an index of all articles published since January 1975 (abstracts 1975-1992, full-text 1993-present). You can search by author, key word, title, type of article, and date. The results will include the citations for the articles plus links to the abstracts of articles published since 1993. For nonsubscribers, time-limited access to single articles and 24-hour site access can also be ordered for a fee through the Internet (<http://www.nejm.org>).

---

## CORRECTION

**Outcomes in Young Adulthood for Very-Low-Birth-Weight Infants**

*To the Editor:* Hack et al. (Jan. 17 issue)<sup>1</sup> report that 20-year-olds who had very low birth weight have a lower rate of risk-taking behavior than their normal-birth-weight peers, and the authors describe this finding as "reassuring." McCormick and Richardson, in their editorial,<sup>2</sup> suggest that the avoidance of risk-taking behavior indicates a special "resilience" in very-low-birth-weight children and their families.

I disagree. As the parent of a very-low-birth-weight adult, the moderator of an Internet list for parents of preterm children, and the author of a book on prematurity, I am in close contact with many families with very-low-birth-weight children. Our children, even when they do not have major neurosensory handicaps, often have cognitive and behavioral deficits that isolate them from both their peers and their peers' risk-taking behavior. Our children's isolation and withdrawal are actually caused by a lack of social and intellectual resilience. As a result, many of us worry that our children will never become fully functioning members of society.

Unfortunately, recent research supports our fears. In a report on a national cohort of prematurely born teens in the Netherlands, Walther et al.<sup>3</sup> estimate that, because of social and cognitive problems, 40 percent of very-low-birth-weight children will never live independently. This Dutch cohort was born only a few years later than the group studied by Hack et al. and has a similar rate of neurosensory impairment (10 percent).

Helen Harrison  
1144 Sterling Ave.  
Berkeley, CA 94708  
helen1144@aol.com

**References**

1. Hack M, Flannery DJ, Schluchter M, Cartar L, Borawski E, Klein N. Outcomes in young adulthood for very-low-birth-weight infants. *N Engl J Med* 2002;346:149-157.
2. McCormick MC, Richardson DK. Premature infants grow up. *N Engl J Med* 2002;346:197-198.
3. Walther FJ, den Ouden AL, Verloove-Vanhorick SP. Looking back in time: outcome of a national cohort of very preterm infants born in the Netherlands in 1983. *Early Hum Dev* 2000;59:175-191.

*To the Editor:* Hack et al. use either a general equivalency diploma or a standard high-school diploma as a measure of academic achievement. The two are not equivalent.<sup>1</sup> If the authors had used only the

latter criterion as a measure of academic success, it is likely that the shortfall in academic achievement among very-low-birth-weight persons would be even more dramatic than that presented. In a nationally representative study,<sup>2</sup> we found that a low-birth-weight child is 74 percent less likely than his or her normal-birth-weight sibling to complete high school by 19 years of age.

In addition, it is not surprising that the less fortunate very-low-birth-weight adults who have chronic disabilities such as blindness, cerebral palsy, or lung disease would be unlikely to be found on the wrong side of the law. Hack and her colleagues state that the relation persisted when they limited their comparison to healthy very-low-birth-weight adults and normal-birth-weight adults. These results should be presented.

Dalton Conley, Ph.D., M.P.A.  
New York University  
New York, NY 10003  
dalton.conley@nyu.edu

Neil G. Bennett, Ph.D.  
City University of New York  
New York, NY 10010

**References**

1. Cameron SV, Heckman JJ. The nonequivalence of high school equivalents. *J Labor Econ* 1993;11:1-47.
2. Conley D, Bennett NG. Is biology destiny? Birth weight and life chances. *Am Sociol Rev* 2000;65:458-467.

*To the Editor:* Hack and colleagues state that very-low-birth-weight babies have a significantly lower mean IQ at 20 years of age than do the members of a control group (87 vs. 92). Although this difference may achieve statistical significance, such a difference is not considered meaningful by those who specialize in assessing cognitive development. Both scores are rated as falling within the average range. IQ scores are not finite measures of a characteristic in a given person; they are merely scores of someone's performance on a given test at a given time and are subject to errors of measurement. In this case, the range of "true" scores results in considerable overlap between the two groups of subjects.

It is known that low-birth-weight babies are at risk for cognitive deficits. What parents of these babies want to know is the nature and extent of this risk. Hack et al. do not emphasize that 120 of the low-birth-weight adults had normal IQs of 85 or higher.

Lillian J. Zach, Ph.D.  
Ferkof Graduate School of Psychology  
Bronx, NY 10461

*To the Editor:* Hack et al. report higher rates of neurosensory impairment among very-low-birth-weight infants. Since premature in-

fants frequently have retinal problems that may have lifelong consequences, I wonder whether the authors were able to identify visual impairment as one of the serious neurosensory problems.

William Tasman, M.D.  
Wills Eye Hospital  
Philadelphia, PA 19107  
wst1@ureach.com

The authors reply:

*To the Editor:* We agree with Harrison that social isolation may have a role in the study subjects' tendency to engage in less risk-taking behavior. However, we do not have information on social relationships.

The estimate of Walther et al. that 40 percent of very-low-birth-weight children will not live independently was based on the responses to questions asked over the telephone of parents of 14-year-old children.<sup>1</sup> We interviewed and tested young adults. Although fewer very-low-birth-weight men than control men were in college, more were working (47 percent vs. 27 percent,  $P < 0.01$ ). Very-low-birth-weight women did not differ significantly from control women in terms of rates of college enrollment or employment. These results indicate that most very-low-birth-weight adults will be able to work and live independently, although men might lag behind in educational attainment.

In response to Conley and Bennett: we performed additional analyses excluding subjects with a general equivalency diploma. The rates of high-school graduation for very-low-birth-weight and normal-birth-weight men were 60 percent and 68 percent, respectively ( $P = 0.28$ ); the rates for women were 77 percent and 84 percent, respectively ( $P = 0.07$ ). When subjects with neurosensory impairment, a subnormal IQ, or both were excluded, the rates of alcohol use for very-low-birth-weight and normal-birth-weight subjects were 69 percent and 84 percent, respectively ( $P = 0.001$ ), and the rates of illicit-drug use were 37 percent and 47 percent, respectively ( $P = 0.02$ ). Fewer men with very low birth weight than with normal birth weight had been in contact with the police for drug-related or alcohol-related offenses (13 percent vs. 29 percent,  $P = 0.008$ ). When we excluded all subjects with chronic conditions (neurosensory, medical, or psychiatric conditions or subnormal IQ), subjects with very low birth weight still had lower rates of alcohol use (68 percent vs. 83 percent,  $P = 0.001$ ) and illicit-drug use (36 percent vs. 49 percent,  $P = 0.009$ ) than normal-birth-weight subjects. Among men, the rates of contact with police for offenses related to drugs or alcohol were 14 percent and 28 percent, respectively ( $P = 0.04$ ).

Tasman asks about visual impairment. Four very-low-birth-weight subjects (1.7 percent) had blindness due to retinopathy of prematurity (bilateral in one subject and unilateral in three).

We agree with Zach that many of the very-low-birth-weight subjects had normal IQs in young adulthood. However, as we noted in the discussion, our results are applicable only to current survivors of neonatal intensive care with birth weights between 1000 g and 1500 g. We

have serious concern about children born during the 1990s weighing less than 1000 g, who may not function well as young adults.<sup>2,3</sup>

In Table 3 of our article, the total number of normal-birth-weight men with postsecondary study should have been 56 rather than 57.

Maureen Hack, M.B., Ch.B.  
Nancy Klein, Ph.D.  
Daniel J. Flannery, Ph.D.  
University Hospitals of Cleveland  
Cleveland, OH 44106  
mxh7@po.cwru.edu

## References

- Walther FJ, den Ouden AL, Verloove-Vanhorick SP. Looking back in time: outcome of a national cohort of very preterm infants born in the Netherlands in 1983. *Early Hum Dev* 2000;59:175-191.
- Hack M, Wilson-Costello D, Friedman H, Taylor GH, Schluchter M, Fanaroff AA. Neurodevelopment and predictors of outcomes of children with birth weights of less than 1000 g: 1992-1995. *Arch Pediatr Adolesc Med* 2000;154:725-731.
- Vohr BR, Wright LL, Dusick AM, et al. Neurodevelopmental and functional outcomes of extremely low birth weight infants in the National Institute of Child Health and Human Development Neonatal Research Network, 1993-1994. *Pediatrics* 2000;105:1216-1226.

The editorialists reply:

*To the Editor:* Ms. Harrison offers an alternative hypothesis for the relative absence of risk-taking behavior observed in the group of young adults studied by Hack et al. Both our somewhat more optimistic hypothesis of resilience and hers of social isolation are testable in follow-up studies of very-low-birth-weight children now approaching adulthood. Her letter underscores the importance of not simply reporting on the outcomes of these vulnerable children, but also exploring the mechanisms that cause them, as we have argued elsewhere.<sup>1</sup> Well-targeted interventions have been demonstrated to effect changes in preschool cognitive and behavioral outcomes in very-low-birth-weight children.<sup>2</sup> Understanding the mechanisms behind other adverse outcomes could lead to the development of strategies for amelioration.

Marie C. McCormick, M.D., Sc.D.  
Harvard School of Public Health  
Boston, MA 02115  
mmccormi@hsph.harvard.edu

Douglas K. Richardson, M.D., M.B.A.  
Beth Israel Deaconess Medical Center  
Boston, MA 02215

## References

1. McCormick MC. Conceptualizing child health status: observations from studies of very premature infants. *Perspect Biol Med* 1999;42:372-386.
2. McCormick MC, McCarton C, Tonascia J, Brooks-Gunn J. Early educational intervention for very low birth weight infants: results from the Infant Health and Development Program. *J Pediatr* 1993;123:527-533.