

Special Article

EVIDENCE OF REAL-WORLD EFFECTIVENESS
OF A TELEPHONE QUITLINE FOR SMOKERSSHU-HONG ZHU, PH.D., CHRISTOPHER M. ANDERSON, B.A., GARY J. TEDESCHI, PH.D., BRADLEY ROSBROOK, M.S.,
CYNTHIA E. JOHNSON, B.A., MICHAEL BYRD, M.A., AND ELSA GUTIÉRREZ-TERRELL, M.A.**ABSTRACT**

Background Telephone services that offer smoking-cessation counseling (quitlines) have proliferated in recent years, encouraged by positive results of clinical trials. The question remains, however, whether those results can be translated into real-world effectiveness.

Methods We embedded a randomized, controlled trial into the ongoing service of the California Smokers' Helpline. Callers were randomly assigned to a treatment group (1973 callers) or a control group (1309 callers). All participants received self-help materials. Those in the treatment group were assigned to receive up to seven counseling sessions; those in the control group could also receive counseling if they called back for it after randomization.

Results Counseling was provided to 72.1 percent of those in the treatment group and 31.6 percent of those in the control group (mean, 3.0 sessions). The rates of abstinence for 1, 3, 6, and 12 months, according to an intention-to-treat analysis, were 23.7 percent, 17.9 percent, 12.8 percent, and 9.1 percent, respectively, for those in the treatment group and 16.5 percent, 12.1 percent, 8.6 percent, and 6.9 percent, respectively, for those in the control group ($P < 0.001$). Analyses factoring out both the subgroup of control subjects who received counseling and the corresponding treatment subgroup indicate that counseling approximately doubled abstinence rates: rates of abstinence for 1, 3, 6, and 12 months were 20.7 percent, 15.9 percent, 11.7 percent, and 7.5 percent, respectively, in the remaining subjects in the treatment group and 9.6 percent, 6.7 percent, 5.2 percent, and 4.1 percent, respectively, in the remaining subjects in the control group ($P < 0.001$). Therefore, the absolute difference in the rate of abstinence for 12 months between the remaining subjects in the treatment and control groups was 3.4 percent. The 12-month abstinence rates for those who made at least one attempt to quit were 23.3 percent in the treatment group and 18.4 percent in the control group ($P < 0.001$).

Conclusions A telephone counseling protocol for smoking cessation, previously proven efficacious, was effective when translated to a real-world setting. Its success supports Public Health Service guidelines calling for greater availability of quitlines. (N Engl J Med 2002;347:1087-93.)

Copyright © 2002 Massachusetts Medical Society.

AMONG services recommended by the U.S. Preventive Services Task Force, tobacco-cessation counseling is ranked in the highest priority category with the lowest delivery rate.¹ The clinical guidelines of the Public Health Service recommend use of the telephone to deliver cessation-counseling services, in part because such "quitlines" have the potential to reach large numbers of smokers.² In recent years, such programs have proliferated. Thirty-three states have established quitlines, and more are preparing to do so. Many other countries have established national quitlines.

The Public Health Service recommendation is based on positive results of clinical trials,²⁻⁵ but the question remains whether quitlines can translate these results into real-world effectiveness. Proven treatments sometimes fail in practice, because translation from clinical trials to service settings may involve changes in the conditions under which the original results were obtained.^{6,7} The staff may differ in skills and enthusiasm, and quality control, a critical element in behavioral interventions, may suffer under the pressure to meet clients' expressed needs. Given the increasing public investment in quitlines, it is therefore important to determine whether such services can maintain the effectiveness found in clinical trials. We examined the real-world effectiveness of the California Smokers' Helpline, a statewide quitline based on a successful clinical trial.⁸ In operation since 1992, the program has been the model for many other quitlines.⁹

The most rigorous way to assess effectiveness in practice would be to conduct a randomized, controlled trial in the context of service operation. However, few smokers calling a fully operational quitline would expect to be assigned to a control group. Furthermore, to impose a control in this setting would compromise ethical standards. These considerations may lead quitline operators to evaluate their services without a randomized design.

From the Department of Family and Preventive Medicine, University of California, San Diego, La Jolla. Address reprint requests to Dr. Zhu at the University of California, San Diego, 9500 Gilman Dr., Mail Code 0905, La Jolla, CA 92093-0905, or at szhu@ucsd.edu.

who did not call back and thus did not receive counseling were designated control subgroup B.

All subjects in the treatment group were immediately assigned to a counselor. Because all participants received the same instructions and group assignment was random, it was inferred that if the treatment group had not received counseling, it would have been divided into two subgroups similar to the two subgroups of the control group: those who would have called back (treatment subgroup A) and those who would not have called back (treatment subgroup B). In Figure 1, the theoretical nature of this division is indicated by a dashed line. Although it is unknown which members of the treatment group would actually have belonged to which subgroup, abstinence rates for treatment subgroup A and treatment subgroup B can nonetheless be derived algebraically (see the Statistical Analysis section). Control subgroup B and treatment subgroup B were equivalent except with respect to service; thus, control subgroup B served as a control for treatment subgroup B in the determination of the effect of counseling.¹⁰

This randomization procedure did not affect the participants' ability to receive treatment, because all smokers, including those in the control group, received the same callback instructions, and those in the control group were not denied counseling if they did call back. Therefore, callers who wanted treatment were not discouraged from participating in the study, and we were able to recruit from the full spectrum of callers. Structurally, this design is similar to Zelen's randomization,¹¹ except that the participants were randomly assigned to groups after consenting to evaluation, not before.

Intervention

The California Smokers' Helpline used a previously tested, structured counseling protocol,¹² which helped to ensure consistency by guiding the content and quality of each session. The first session focused on quitting history, motivation, self-efficacy, social support, and planning in advance of the quitting date. Sessions after the quitting date (up to six sessions) emphasized relapse prevention by reviewing and revising the plan and promoting adoption of the self-image of a nonsmoker. All sessions were conducted proactively (i.e., the counselor called the smoker) and occurred within three months after the initial contact. The counseling protocol and quality-assurance methods used in the study are described in detail elsewhere.¹² Twenty English-speaking counselors, six of whom also spoke Spanish, provided treatment. All had at least a bachelor's degree, and six had master's degrees. All received 60 hours of training, regardless of their background.

Follow-up Assessment

A separate staff of evaluators interviewed the participants 2, 4, 7, and 13 months after the initial contact. This schedule was chosen over the more common 12-month schedule because previous experience indicated a lag between the time smokers call for service and the time they actually try to quit. In order to assess prolonged abstinence (at 12 months), an extra month was allowed. All follow-up was conducted by telephone. Smoking status was based on self-report, since biochemical tests are considered uninformative in low-intensity intervention studies such as this.^{13,14}

Statistical Analysis

The effects of intervention were first tested by directly comparing the overall treatment and control groups. To provide a better numerical estimate of the effect of counseling — that is, the difference between treatment subgroup B and control subgroup B — we then factored out the control subgroup that received counseling (control subgroup A), along with its counterpart in the treatment group (treatment subgroup A). Because of the randomization procedure and the fact that treatment subgroup A and control subgroup A both received the same intervention, they were assumed

to have the same quitting rate. The quitting rate in treatment subgroup B (Q_{TB}) was estimated by the following formula: $Q_{TB} = (Q_T - w_{CA}Q_{CA}) \div w_{CB}$, where Q_T is the quitting rate in the treatment group, Q_{CA} is the quitting rate in control subgroup A, and w_{CA} and w_{CB} are the relative proportions of participants in control subgroup A and control subgroup B, respectively ($w_{CA} + w_{CB} = 1$).¹⁰

Multiple-outcome measures were used in the analysis, according to the recommendations of the Society for Research on Nicotine and Tobacco for assessing treatment effects.¹⁵ The primary measure of the overall intervention effect was prolonged abstinence on the basis of an intention-to-treat approach. Four periods of abstinence commensurate with the evaluation schedule were assessed: 1, 3, 6, and 12 months. Life-table estimates were computed for all four intervals, and a log-rank test was used.¹⁶ In the primary analysis, those lost to follow-up were considered to have continued smoking.

Additional measures were the percentages of smokers who quit for at least 24 hours (i.e., made a serious attempt to quit) in the first three months and the probability of relapse over time for these attempts, with relapse defined as two consecutive days of smoking.⁸ Because three months was the longest period during which a participant was in contact with a counselor, quitting attempts occurring after three months were not attributed to counseling. Abstinence curves were obtained by the Kaplan–Meier product-limit method.¹⁶ Comparison of these curves gives a clear picture of the effect of counseling on the prevention of relapse, as distinct from its effect on attempts to quit. For these secondary analyses, participants for whom there were no follow-up data were excluded, because even if we assumed they were smokers, it was not known whether they had attempted to quit and relapsed or simply never attempted to quit.

RESULTS

Base-Line Characteristics

No significant difference in base-line characteristics was found between the two study groups (Table 1). The characteristics of the study participants were similar to those of callers who chose counseling but did not participate in the study.

Percentages of Subjects Receiving Counseling

Of the 1309 subjects in the control group, 463 (35.4 percent) called back for counseling. These subjects (control subgroup A) were assigned to counselors, as were all 1973 subjects in the treatment group. The counselors attempted to contact everyone assigned to them, counseling 31.6 percent of those in the control group (89.4 percent of control subgroup A) and 72.1 percent of those in the treatment group. Thus, those in the treatment group received counseling at a rate 40.5 percentage points higher than those in the control group. Counseled participants received an average of 3.0 counseling sessions (3.0 and 2.9 for the control and treatment groups, respectively).

Rates of Prolonged Abstinence

As shown in Table 2, 16.5 percent of the control subjects and 23.7 percent of the treatment subjects quit smoking for at least one month. The rates of prolonged abstinence decreased over time in both

TABLE 1. BASE-LINE CHARACTERISTICS OF SUBJECTS IN THE TWO STUDY GROUPS AND OF NONPARTICIPANTS.*

CHARACTERISTIC	CONTROL GROUP (N=1309)	TREATMENT GROUP (N=1973)	NON-PARTICIPANTS (N=3501)†
Sex (%)			
Male	42.4±2.7	44.7±2.7	45.1±1.7
Female	57.6±2.6	55.3±2.7	54.9±1.7
Age (yr)	37.9±0.6	38.6±0.6	39.1±0.4
Education (%)			
High school or less	50.8±2.7	47.6±2.7	47.0±1.7
Some college	34.2±2.6	38.2±2.6	37.3±1.6
College graduate	15.0±1.9	14.2±1.8	15.7±1.2
Race or ethnic group (%)			
White	61.9±2.6	62.9±2.6	65.4±1.6
Hispanic	22.0±2.2	20.9±2.2	17.7±1.3
Black	10.1±1.6	10.3±1.7	9.7±1.0
Asian	1.5±0.7	1.8±0.7	1.8±0.4
American Indian	2.6±0.9	2.3±0.8	3.2±0.6
Other	1.9±0.7	1.7±0.7	2.2±0.5
No. of cigarettes smoked daily	20.1±0.6	21.0±0.7	21.2±0.4
No. of previous attempts to quit			
0	15.7±2.0	14.0±1.9	14.7±1.2
1-2	38.6±2.6	37.1±2.6	39.2±1.6
>2	45.7±2.7	48.9±2.7	46.1±1.7
Residing with other smokers (%)	42.1±2.7	40.3±2.7	42.2±1.6

*Values are means with 95 percent confidence intervals. Because of rounding, not all percentages add to 100.

†Nonparticipants were adult smokers who chose counseling but did not participate in the randomized study.

groups: 6.9 percent of the control subjects and 9.1 percent of the treatment subjects quit smoking for at least one year ($P<0.001$ by the log-rank test).

The rates of prolonged abstinence were much higher in control subgroup A than in control subgroup B (Table 2). Assuming that treatment subgroup A and control subgroup A were equivalent allowed the success rates in treatment subgroup B to be derived algebraically. The rates of abstinence for 1, 3, 6, and 12 months in treatment subgroup B were approximately twice those in control subgroup B ($P<0.001$).

Rates of Quitting Attempts and Probability of Relapse

Secondary analyses examined the effects of intervention on attempts to quit during the first three months and on the probability of relapse after those attempts. As shown in Table 3, the rate of quitting attempts was higher in the treatment group than in the control group (62.9 percent vs. 56.9 percent, $P<0.001$). By a wider margin, subjects in treatment subgroup B were more likely to attempt to quit than those in control subgroup B (57.9 percent vs. 48.6 percent, $P<0.001$). Not surprisingly, those in control subgroup A were much more likely to try to quit than those in control subgroup B (71.9 percent vs. 48.6 percent).

Abstinence curves for these attempts to quit are given in Figure 2. Among smokers who made an attempt to quit, those in treatment subgroup A and

TABLE 2. RATES OF PROLONGED ABSTINENCE ACCORDING TO THE INTENTION-TO-TREAT ANALYSIS.*

STUDY GROUP	LENGTH OF ABSTINENCE†			
	1 MO	3 MO	6 MO	12 MO
	percent of subjects			
Control (n=1309)	16.5	12.1	8.6	6.9
Treatment (n=1973)	23.7	17.9	12.8	9.1
Subgroup analysis				
Control				
Control subgroup B (self-help, n=846)	9.6	6.7	5.2	4.1
Control subgroup A (counseling, n=463)	29.2	21.6	14.9	11.9
Treatment‡				
Treatment subgroup B (counseling, n=1275)	20.7	15.9	11.7	7.5
Treatment subgroup A (counseling, n=698)	29.2	21.6	14.9	11.9

*All participants who were lost to follow-up were considered current smokers. The percentages of subjects with follow-up data at 1, 3, 6, and 12 months were 90.6, 86.1, 80.7, and 69.9 percent for control subjects and 89.1, 85.7, 80.3, and 71.1 percent for treatment-group subjects, respectively (P not significant).

†Log-rank tests comparing control and treatment groups and comparing control subgroup B and treatment subgroup B showed significant differences for all four intervals ($P<0.001$).

‡These percentages were not directly measured. Estimates for treatment subgroup B (Q_{TB}) were obtained by the equation $Q_{TB} = (Q_T - w_{CA}Q_{CA}) \div w_{CB}$, where Q_T is the quitting rate in the treatment group, Q_{CA} is the quitting rate in control subgroup A, and w_{CA} and w_{CB} are the proportions of subjects in control subgroups A and B, respectively (see Zhu¹⁰ for details). The quitting rates were assumed to be the same in treatment subgroup A and control subgroup A.

TABLE 3. RATES OF ATTEMPTS TO QUIT SMOKING FOR 24 HOURS WITHIN THE FIRST 3 MONTHS AND 12-MONTH ABSTINENCE RATES.*

STUDY GROUP	ATTEMPTS TO QUIT WITHIN 3 Mo		12-Mo ABSTINENCE RATE AFTER ATTEMPTING TO QUIT	
	%	P VALUE	%	P VALUE
Control (n=1190)	56.9	<0.001	23.3	0.04
Treatment (n=1761)	62.9		25.8	
Subgroup analysis				
Control subgroup B (self-help)†	48.6	<0.001	18.4	<0.001
Treatment subgroup B (counseling)	57.9		23.3	

*Participants for whom the California Smokers' Helpline had no follow-up data were excluded from these analyses.

†Estimates for treatment subgroup B (Q_{TB}) were obtained by the equation $Q_{TB} = (Q_T - w_{CA}Q_{CA}) \div w_{CB}$, where Q_T is the quitting rate in the treatment group, Q_{CA} is the quitting rate in control subgroup A, and w_{CA} and w_{CB} are the proportions of subjects in control subgroups A and B, respectively (see Zhu¹⁰ for details).

control subgroup A, which received counseling and were assumed to be equivalent, had the highest rate of long-term abstinence; those in treatment subgroup B, which received counseling, had a higher rate of long-term abstinence than those in control subgroup

B, which received only self-help materials ($P < 0.001$).

The 12-month end points for these groups are shown in Table 3. The 12-month abstinence rate was higher in the treatment group than in the control group (25.8 percent vs. 23.3 percent, $P = 0.04$). Again by a wider margin, the 12-month abstinence rate was higher in treatment subgroup B than in control subgroup B (23.3 percent vs. 18.4 percent, $P < 0.001$). Treatment subgroup A, which was assumed to be equivalent to control subgroup A, had an even higher abstinence rate (29.4 percent, $P < 0.001$).

DISCUSSION

In this study, a telephone counseling protocol for smoking cessation that had previously proved efficacious⁸ maintained its effectiveness when translated to a real-world setting. The treatment effect in this study was obtained under demanding conditions in which a substantial portion of smokers randomly assigned to a control group (31.6 percent) received the same counseling service as those in treatment. Moreover, only 72.1 percent of smokers randomly assigned to the treatment group received counseling, as often happens in real-world settings where participants drop out before receiving treatment.

There was a difference of only 40.5 percentage




Figure 2. Relapses in the Treatment and Control Subgroups.

Treatment subgroup A (counseling) and control subgroup A (counseling), which were assumed to be equivalent, were the most successful subgroups and appear as a single curve. Control subgroup B (self-help) was the least successful. Between these is a curve representing treatment subgroup B (counseling). (The curve is smooth to indicate that it was derived algebraically.) Quitting attempts made by subjects in treatment subgroup B were more likely to lead to long-term abstinence than those made by subjects in control subgroup B ($P < 0.001$).

points between the two groups in the rates of counseling. As a result, the absolute effect of the counseling intervention was limited. The absolute difference in 12-month abstinence rates between the counseling group (treatment subgroup B) and the self-help group (control subgroup B) was 3.4 percent. However, as Table 2 shows, when the control subgroup that received counseling and the corresponding treatment subgroup were factored out, the prolonged abstinence rates in treatment subgroup B were roughly twice those in control subgroup B for all four periods of abstinence.

Further analysis indicates that the intervention achieved its effect through two mechanisms: by increasing the percentage of smokers making attempts to quit and by reducing the probability of relapse (as can be seen in the comparison of treatment subgroup B with control subgroup B in Table 3). Moreover, it seems likely that the smokers in treatment subgroup B had greater ambivalence than those in treatment subgroup A. They would not have used counseling if counselors had not proactively called them. The higher success rate in treatment subgroup B than in control subgroup B therefore provides empirical support for the effectiveness of proactive counseling.^{2,5,8,17}

This study has several limitations. Although its design succeeded in achieving random assignment in a public health setting, it could not accurately assess the extent to which counseling contributed to success in control subgroup A and treatment subgroup A, because there was no randomized control specifically for these subgroups. Smokers in these subgroups followed the instructions to call back to start the counseling, and not surprisingly,^{18,19} they were the most successful in quitting. Because all three subgroups received the same counseling, it is unknown whether counseling doubled the quitting rates of subjects in these subgroups, just as it did for those in treatment subgroup B.

The estimation of abstinence rates for treatment subgroup B assumes equality between treatment subgroup A and control subgroup A, given the random assignment of participants to either treatment or control. Although Table 1 shows that randomization produced equivalent groups at base line, a direct test of the equality of treatment subgroup A and control subgroup A was not possible, because it was not known which members of the treatment group belonged to which subgroup. However, we found a significant difference in abstinence rates between the overall treatment and control groups, a comparison that was not dependent on any assumptions about the equality of treatment subgroup A and control subgroup A.^{10,11}

Although this study shows that the quitline's counseling was effective, its direct clinical effect is limited by the fact that most smokers in the program had re-

lapsed by the end of 12 months. Nevertheless, the doubling effect of counseling on success rates compares favorably with the clinical-trial results summarized by the Public Health Service in its recommendation for telephone counseling.² Moreover, quitlines are often an integral part of statewide antitobacco media campaigns, so they may have an effect on public health beyond that of the counseling alone.²⁰ Perhaps for that reason, many states allocate a large share of their smoking-cessation funds to support widely promoted, centrally operated quitlines rather than traditional cessation clinics. A detailed discussion of the cost effectiveness of quitlines relative to that of clinics is beyond the scope of this paper, but recent data suggest that smokers are four times as likely to use quitlines as face-to-face clinics, given the same promotional effort, so the savings in promotion alone could be substantial.²¹

Because we were able to embed a randomized trial within the ongoing service of a large statewide quitline, this study provides strong evidence that telephone counseling conducted in a real-world setting can be an effective means of helping smokers quit. This evidence supports the Public Health Service guidelines calling for greater availability of telephone counseling services for smoking cessation.

Supported by funds received from the Tobacco Tax Health Protection Act of 1988, Proposition 99, under a grant (96-27049) from the California Department of Health Services.

REFERENCES

1. Coffield AB, Maciosek MV, McGinnis JM, et al. Priorities among recommended clinical preventive services. *Am J Prev Med* 2001;21:1-9.
2. Fiore MC, Bailey WC, Cohen SJ, et al. Treating tobacco use and dependence. Clinical practice guideline. Rockville, Md.: Public Health Service, June 2000.
3. Lichtenstein E, Glasgow RE, Lando HA, Ossip-Klein DJ, Boles SM. Telephone counseling for smoking cessation: rationales and meta-analytic review of evidence. *Health Educ Res* 1996;11:243-57.
4. Stead LF, Lancaster T. Telephone counselling for smoking cessation (Cochrane review). In: *The Cochrane library*, 1. Oxford, England: Update Software, 2002 (data base).
5. Hopkins DP, Briss PA, Ricard CJ, et al. Reviews of evidence regarding interventions to reduce tobacco use and exposure to environmental tobacco smoke. *Am J Prev Med* 2001;20:Suppl:16-66.
6. Peterson AL, Halstead TS. Group cognitive behavior therapy for depression in a community setting: a clinical replication series. *Behav Ther* 1998;29:3-18.
7. Stevens VJ, Glasgow RE, Hollis JF, Mount K. Implementation and effectiveness of a brief smoking-cessation intervention for hospital patients. *Med Care* 2000;38:451-9.
8. Zhu S-H, Stretch V, Balabanis M, Rosbrook B, Sadler G, Pierce JP. Telephone counseling for smoking cessation: effects of single-session and multiple session interventions. *J Consult Clin Psychol* 1996;64:202-11.
9. Zhu S-H, Anderson CM, Johnson CE, Tedeschi G, Roeseler A. A centralized telephone service for tobacco cessation: the California experience. *Tob Control* 2000;9:Suppl 2:II-48-II-55.
10. Zhu S-H. A method to obtain a randomized control group where it seems impossible: a case study in program evaluation. *Eval Rev* 1999;23:363-77.
11. Zelen M. A new design for randomized clinical trials. *N Engl J Med* 1979;300:1242-5.
12. Zhu S-H, Tedeschi G, Anderson CM, Pierce JP. Telephone counseling for smoking cessation: what's in a call? *J Couns Dev* 1996;75:93-102.
13. Velicer WF, Prochaska JP, Rossi JS, Snow MG. Assessing outcome in smoking cessation studies. *Psychol Bull* 1992;111:23-41.

14. Benowitz NL, Jacob P III, Ahijevych K, et al. Biochemical verification of tobacco use and cessation. *Nicotine Tob Res* 2002;4:149-59.
15. Hughes JR, Keely JP, Niaura RS, Ossip-Klein DJ, Richmond RL, Swan GE. Measures of abstinence from tobacco in clinical trials. *Nicotine Tob Res* (in press).
16. Kalbfleisch JD, Prentice RL. *The statistical analysis of failure time data*. New York: John Wiley, 1980.
17. Miller WR, Rollnick S. *Motivational interviewing: preparing people to change addictive behavior*. New York: Guilford Press, 1991.
18. Horwitz RI, Horwitz SM. Adherence to treatment and health outcomes. *Arch Intern Med* 1993;153:1863-8.
19. McDermott MM, Schmitt B, Wallner E. Impact of medication nonadherence on coronary heart disease outcomes: a critical review. *Arch Intern Med* 1997;157:1921-9.
20. Wakefield M, Borland R. Saved by the bell: the role of telephone help-line services in the context of mass-media anti-smoking campaigns. *Tob Control* 2000;9:117-9.
21. McAfee T, Sofian NS, Wilson J, Hindmarsh M. The role of tobacco intervention in population-based health care: a case study. *Am J Prev Med* 1998;14:46-52.

Copyright © 2002 Massachusetts Medical Society.