



This Week in the Journal

JULY 15, 2004

CORRESPONDENCE

- 295 C-Reactive Protein and Coronary Heart Disease
- 298 Childhood Vaccination and Type 1 Diabetes
- 298 Bioethics, Science, and Politics
- 300 Management of Cirrhosis and Ascites
- 301 Pediatric Palliative Care
- 302 Blue Cohosh and Perinatal Stroke

BOOK REVIEWS

- 304 The Genome War: How Craig Venter Tried to Capture the Code of Life and Save the World
- 305 Flesh in the Age of Reason

CONTINUING MEDICAL EDUCATION

- 309 Treatment of Deep-Vein Thrombosis
- 310 Timing and Magnitude of Increases in Levothyroxine Requirements during Pregnancy in Women with Hypothyroidism
- 311 Single-Dose Perinatal Nevirapine plus Standard Zidovudine to Prevent Mother-to-Child Transmission of HIV-1 in Thailand

Next Week in the Journal

JULY 22, 2004

The High Cost of New Cancer Therapies

Deborah Schrag

ORIGINAL ARTICLE

Nevirapine to Prevent Mother-to-Child Transmission of HIV-1

In a randomized, controlled trial in Thailand, pregnant women infected with the human immunodeficiency virus (HIV) type 1 received zidovudine treatment plus a single dose of either nevirapine or placebo. The rate of transmission of HIV to newborns was 2.8 percent in the nevirapine group and 6.3 percent in the placebo group. The rate was 1.9 percent in a group in which a single dose of nevirapine was also administered to the infants shortly after delivery.

When added to zidovudine prophylaxis, a single dose of nevirapine is highly effective in reducing mother-to-infant transmission of HIV.

SEE P. 217; EDITORIAL, P. 289

ORIGINAL ARTICLE

Subsequent Maternal Effects of Intrapartum Exposure to Nevirapine

After participating in a placebo-controlled trial of antiviral regimens to reduce mother-to-child transmission of the human immunodeficiency virus (HIV) in Thailand, some of the women began taking a nevirapine-containing regimen. After six months, the rate of viral suppression was lower among the mothers who had received intrapartum nevirapine than among those who had not received it (49 percent vs. 68 percent, $P=0.03$).

These observational data suggest that intrapartum exposure to a single dose of nevirapine may lead to selection for resistance mutations that can adversely affect a woman's subsequent antiviral treatment.

SEE P. 229; EDITORIAL, P. 289

ORIGINAL ARTICLE

Levothyroxine Requirements in Women with Hypothyroidism during Pregnancy

Despite knowledge that women with hypothyroidism should increase their usual levothyroxine dose during pregnancy, biochemical hypothyroidism still occurs. This study found that the levothyroxine requirement increased by a mean of 48 percent during the first half of pregnancy and that monitoring and dose adjustments were required until delivery.

Given the importance of maternal euthyroidism for normal fetal cognitive development, women with hypothyroidism should increase their levothyroxine dose by approximately 30 percent once pregnancy is confirmed.

SEE P. 241; EDITORIAL, P. 292

ORIGINAL ARTICLE

Lymphoma-Specific Genetic Aberrations in Microvascular Endothelial Cells in B-Cell Lymphomas

Using combined immunohistochemical and fluorescence in situ hybridization techniques, these investigators found chromosomal abnormalities typical of B-cell lymphomas not only in the lymphoma cells but also in the microvascular endothelial cells in the lymphoma.

These findings link lymphoma cells to angiogenesis. The discovery of how this link forms will be a major step forward in understanding the origins and growth of lymphomas.

SEE P. 250; PERSPECTIVE, P. 215

CLINICAL PRACTICE

Venous Thromboembolism

A 52-year-old woman with no history of venous thromboembolism presents with a four-day history of discomfort in her left calf. Proximal deep-vein thrombosis is diagnosed by compression ultrasonography. How should her case be managed?


SEE P. 268

ORIGINAL ARTICLE

Risk of Arterial Thrombosis in Patients with Systemic Lupus Erythematosus

Arterial thrombosis is an important complication of systemic lupus erythematosus. These investigators found that the presence of variant alleles of mannose-binding lectin (MBL), a serum protein involved in innate immune defense, increases the risk of arterial thrombosis, especially myocardial infarction, in patients with lupus. The results suggest that the common allelic forms of mannose-binding lectin may have a role in protecting against arterial thrombotic events.

SEE P. 260



CASE RECORDS OF THE MASSACHUSETTS GENERAL HOSPITAL

A Woman with a Pericardial Effusion

A woman sought medical attention because of chest pain and shortness of breath. Physical examination and echocardiographic studies showed evidence of a pericardial effusion and cardiac tamponade. She was admitted to the hospital, and the effusion was drained, yielding fluid that was negative on culture and cytologic analysis. She had traveled to Kenya; a tuberculin skin test five months after the trip had been negative, but a test during hospitalization was positive.

SEE P. 279