

paraganglioma syndrome 1; in the extent of multifocal tumors in the group with MEN-2, the group with von Hippel–Lindau disease, and the group with paraganglioma syndrome 1; and in the extent of extraadrenal tumors in the group with MEN-2, the group with von Hippel–Lindau disease, the group with paraganglioma syndrome 1, and the group with paraganglioma syndrome 4.

Patients with neurofibromatosis type 1 had a relatively high (but not significant) prevalence of malignant disease (12 percent), second only to that among patients with paraganglioma syndrome 4 who had a germ-line mutation in the *SDHB* gene (24 percent). Taken together, 33 percent of all symptomatic patients with pheochromocytoma in the registry carried germ-line mutations in one of the five genes, including *NF1*.

Dr. Neumann is supported by a grant (NE 571/5-3) from the Deutsche Forschungsgemeinschaft, a grant (70-3313-ne1) from the Deutsche Krebshilfe, and a grant (LSHC-CT-2005-518200) from the European Union.

Birke Bausch, M.D.
Wiktor Borozdin, Ph.D.
Hartmut P.H. Neumann, M.D.

University of Freiburg
79106 Freiburg, Germany
neumann@med1.ukl.uni-freiburg.de

for the European-American Pheochromocytoma Study Group

1. Riccardi VM. Von Recklinghausen neurofibromatosis. *N Engl J Med* 1981;305:1617-27.
2. Dluhy RG. Pheochromocytoma — death of an axiom. *N Engl J Med* 2002;346:1486-8.
3. Luijten M, Wang Y, Smith BT, et al. Mechanism of spreading of the highly related neurofibromatosis type 1 (*NF1*) pseudogenes on chromosomes 2, 14 and 22. *Eur J Hum Genet* 2000; 8:209-14.
4. Neumann HP, Pawlu C, Peczkowska M, et al. Distinct clinical features of paraganglioma syndromes associated with *SDHB* and *SDHD* gene mutations. *JAMA* 2004;292:943-51. [Erratum, *JAMA* 2004;292:1686.]
5. Borozdin W, Boehm D, Leipoldt M, et al. *SALL4* deletions are a common cause of Okiihiro and acro-renal-ocular syndromes and confirm haploinsufficiency as the pathogenic mechanism. *J Med Genet* 2004;41(9):e113.

Fatal Infection with Influenza A (H5N1) Virus in China

TO THE EDITOR: A 24-year-old man had pneumonia and respiratory distress in November 2003 and died four days after being hospitalized. Because the clinical manifestations were consistent with those of the severe acute respiratory syndrome (SARS) and occurred when sporadic cases of SARS were described in southern China,¹ serum and lung tissue from the patient, as well as fluid aspirated from his chest, were examined for SARS coronavirus with the use of indirect fluorescence antibody tests and the reverse-transcriptase polymerase chain reaction (RT-PCR). All tests were negative for SARS.

A virus was isolated from the lung tissue in Vero-cell cultures and was characteristic of influenza A virus on electron microscopy. A serum sample collected on day 8 of the patient's illness was positive for IgM antibody against the isolated virus. Fragments of both the influenza A virus matrix gene (*M*) and the H5-subtype hemagglutinin gene (*HA*) were amplified from the infected Vero cells with the use of RT-PCR assay.^{2,3} The nucleotide sequences of the fragments were identical to those amplified from the stored specimens of the patient's serum, chest fluid, and lung tissue.

The genomic sequence of the virus (A/Beijing/01/2003) was determined, and its eight segments were genetically related most closely to corresponding sequences of influenza A (H5N1) viruses that had been isolated from chickens in various regions of China in 2004. The segments of the polymerase basic protein 1 gene (*PB*) and the non-structural gene (*NS*) were most closely related to those from Guangdong Province (in southeastern China), with 99 percent identity. The segments of the polymerase basic protein 2 gene (*PB2*) and *HA* gene were closest to those from Jilin Province (in northeastern China), with 99 percent and 97 percent identity, respectively. The segments of the neuraminidase gene (*NA*), nucleoprotein gene (*NP*), and *M* gene were closest to those from Hubei Province (in mideastern China), with 98 percent, 98 percent, and 99 percent identity, respectively, and the polymerase acidic protein gene (*PA*) segment was closest to that from Japan, with 99 percent identity.

These findings suggest that influenza A/Beijing/01/2003 may be a mixed virus. Phylogenetic analyses of the *HA* and *NA* genes of the representative influenza A (H5N1) strains have revealed that the viruses isolated from patients in Thailand

and Vietnam in 2004 and 2005 belong to the same clade, and those obtained from patients in Hong Kong in 1997 and 1998 are from another clade (Fig. 1 in the Supplementary Appendix, available with the full text of this letter at www.nejm.org). A sample of virus obtained from a patient in Hong Kong in 2003 seems to represent a transitional genotype, of which the HA gene sequence was close to the cluster from southeastern Asia (Fig. 1A in the Supplementary Appendix), whereas the NA gene sequence was close to that of the cluster from Hong Kong in 1997 and 1998 (Fig. 1B in the Supplementary Appendix). Phylogenetic analyses of the HA or NA gene indicated that the influenza A/Beijing/01/2003 strain was genetically distant from viruses previously isolated from humans, although it appears to have originated from a lineage similar to the influenza A/goose/Guangdong/1/96 (Gs/GD) lineage.⁴

These findings have important implications for selecting viruses for the development of vaccines to prevent infection in humans. The genetic distance between the isolate reported and the strain currently proposed for vaccine development (A/Vietnam/1203/2004)⁵ implies that viruses from different regions may need to be considered in the development of an effective vaccine against influenza A virus.

Supported by grants from the National Task Force of China (2004BA519A62 and 2004BA519A71) and the 973 High-Tech Projects Plan (2005CB52300015).

Qing-Yu Zhu, M.D.

State Key Laboratory of Pathogens and Biosecurity
Beijing 100071, China

E-De Qin, M.D.

Beijing Institute of Microbiology and Epidemiology
Beijing 100071, China

Wei Wang, M.D.

309th Hospital of the People's Liberation Army
Beijing 100091, China

Jun Yu, Ph.D.

Beijing Genomics Institute
Beijing 101300, China

Bo-Hua Liu, Ph.D.

State Key Laboratory of Pathogens and Biosecurity
Beijing 100071, China

Yi Hu, Ph.D.

Beijing Institute of Microbiology and Epidemiology
Beijing 100071, China

Jian-Fei Hu, Ph.D.

Beijing Genomics Institute
Beijing 101300, China

Wu-Chun Cao, M.D., Ph.D.

State Key Laboratory of Pathogens and Biosecurity
Beijing 100071, China
caowc@nic.bmi.ac.cn

1. Liang G, Chen Q, Xu J, et al. Laboratory diagnosis of four recent sporadic cases of community-acquired SARS, Guangdong Province, China. *Emerg Infect Dis* 2004;10:1774-81.

2. Poddar SK. Influenza virus types and subtypes detection by single step single tube multiplex reverse transcription-polymerase chain reaction (RT-PCR) and agarose gel electrophoresis. *J Virol Methods* 2002;99:63-70.

3. Lee MS, Chang PC, Shien JH, Cheng MC, Shieh HK. Identification and subtyping of avian influenza viruses by reverse transcription-PCR. *J Virol Methods* 2001;97:13-22.

4. Chen H, Smith GJD, Li KS, et al. Establishment of multiple sublineages of H5N1 influenza virus in Asia: implications for pandemic control. *Proc Natl Acad Sci U S A* 2006;103:2845-50.

5. Treanor JJ, Campbell JD, Zangwill KM, Rowe T, Wolff M. Safety and immunogenicity of an inactivated subvirion influenza A (H5N1) vaccine. *N Engl J Med* 2006;354:1343-51.

Correspondence Copyright © 2006 Massachusetts Medical Society.

INSTRUCTIONS FOR LETTERS TO THE EDITOR

Letters to the Editor are considered for publication, subject to editing and abridgment, provided they do not contain material that has been submitted or published elsewhere. Please note the following: •Letters in reference to a *Journal* article must not exceed 175 words (excluding references) and must be received within three weeks after publication of the article. Letters not related to a *Journal* article must not exceed 400 words. All letters must be submitted over the Internet at <http://authors.nejm.org>. •A letter can have no more than five references and one figure or table. •A letter can be signed by no more than three authors. •Financial associations or other possible conflicts of interest must be disclosed. (Such disclosures will be published with the letters. For authors of *Journal* articles who are responding to letters, this information appears in the original articles.) •Include your full mailing address, telephone number, fax number, and e-mail address with your letter.

Our Web site: <http://authors.nejm.org>

We cannot acknowledge receipt of your letter, but we will notify you when we have made a decision about publication. Letters that do not adhere to these instructions will not be considered. Rejected letters and figures will not be returned. We are unable to provide prepublication proofs. Submission of a letter constitutes permission for the Massachusetts Medical Society, its licensees, and its assignees to use it in the *Journal's* various print and electronic publications and in collections, revisions, and any other form or medium.