

# THIS WEEK in the JOURNAL

## ORIGINAL ARTICLE

### Ranibizumab and Age-Related Macular Degeneration

Previous studies have suggested that targeting intravitreal vascular endothelial growth factor A (VEGF-A) counters choroidal neovascularization and hence age-related macular degeneration. A double-blind, placebo-controlled trial of ranibizumab, which neutralizes all isoforms of VEGF-A, for treatment of minimally classic and occult age-related macular degeneration retarded the progression of the disease and improved visual acuity in some patients.

SEE P. 1419; PERSPECTIVE, P. 1409;  
EDITORIAL, P. 1493; CME, P. 1515

## ORIGINAL ARTICLE

### Ranibizumab versus Verteporfin in Age-Related Macular Degeneration

Previous studies have implicated intravitreal vascular endothelial growth factor A (VEGF-A) as a target for countering neovascularization and, therefore, age-related macular degeneration. This double-blind, controlled trial comparing ranibizumab, which neutralizes all isoforms of VEGF-A, with photodynamic therapy with verteporfin showed that ranibizumab was better able to retard the progression of predominantly classic neovascular age-related macular degeneration.

SEE P. 1432; PERSPECTIVE, P. 1409;  
EDITORIAL, P. 1493

## ORIGINAL ARTICLE

### Immunotherapy with a Ragweed–Toll-Like Receptor 9 Agonist Vaccine for Allergic Rhinitis

In standard immunotherapy, small amounts of allergen are injected to induce a state of clinical tolerance of allergen reexposure. In this trial, the investigators studied immunotherapy in which a ragweed allergen was conjugated to an oligonucleotide. The vaccine did not reduce the albumin level in nasal-lavage fluid (the primary end point) but did have a positive effect on an array of secondary end points, suggesting that this approach merits further study in larger, structured trials.

SEE P. 1445

## ORIGINAL ARTICLE

### Lenalidomide in the Myelodysplastic Syndrome with Chromosome 5q Deletion

Anemia is a principal feature of the myelodysplastic syndrome with deletion of chromosome 5q; most patients require transfusions, and iron overload develops as a consequence. In this phase 2 trial, lenalidomide reduced dependence on transfusions in most patients, and two thirds of them could stop transfusions altogether. The drug, an analogue of thalidomide, also reversed chromosomal and cytologic abnormalities in many patients.

SEE P. 1456

## CLINICAL PRACTICE

### Prevention of Meningococcal Disease

A previously healthy 18-year-old college freshman presented at an emergency department with acute onset of fever and headache. He was listless, febrile, and hypotensive. Petechiae were noted over his thorax. Meningococcemia was suspected (and subsequently confirmed by blood cultures positive for *Neisseria meningitidis* serogroup C). Despite prompt administration of antibiotics and other support, the patient's illness was fulminant and he died 12 hours after the onset of symptoms. Should he have previously received meningococcal vaccine, and what measures should be taken to protect his close contacts and his community?

SEE P. 1466; CME, P. 1514

## MECHANISMS OF DISEASE

### Age-Related Macular Degeneration

This review describes the mechanisms entailed in the leading cause of blindness in the United States.

SEE P. 1474; PERSPECTIVE, P. 1409;  
EDITORIAL, P. 1493

## CLINICAL PROBLEM-SOLVING

### Lost in Transcription

Urinary urgency and fever developed in a 55-year-old, bedridden woman with multiple sclerosis, a long-term indwelling Foley catheter, and multiple prior urinary tract infections. The patient had recently been transferred from an assisted-living facility to a skilled-nursing facility because of progressive disability. The laboratory evaluation revealed pancytopenia.

SEE P. 1487; CME, P. 1513